

Helsinki University Central Hospital  
Department of Otorhinolaryngology  
University of Helsinki, Finland

# **PAEDIATRIC OTORHINOLARYNGOLOGICAL FINDINGS IN LUANDA, ANGOLA**

**Anni Taipale**

ACADEMIC DISSERTATION

To be publicly discussed  
by permission of the Faculty of Medicine of the University of Helsinki,  
in the Auditorium 2 of Biomedicum, on October 11, 2013 at 12 noon.

Helsinki 2013

Supervised by	Professor Anne Pitkäranta, MD, PhD Helsinki University Central Hospital Department of Otorhinolaryngology University of Helsinki
Reviewed by	Docent Juha-Pekka Vasama, MD, PhD Tampere University Central Hospital Department of Otorhinolaryngology University of Tampere  Docent Aino Ruohola, MD, PhD Turku University Hospital Department of Paediatrics University of Turku
Official opponent	Professor Malou Hultcrantz, MD, PhD Karolinska University Hospital Ear, Nose and Throat Department Karolinska Institute Stockholm, Sweden

ISBN 978-952-10-9181-0 (nid.)  
ISBN 978-952-10-9182-7 (PDF)  
<http://ethesis.helsinki.fi/>  
Unigrafia Oy, Yliopistopaino  
Helsinki 2013

# 1. CONTENTS

<b>1. Abstract .....</b>	<b>7</b>
<b>Yhteenveto.....</b>	<b>9</b>
 <b>2. Original publications .....</b>	 <b>11</b>
 <b>3. Abbreviations .....</b>	 <b>12</b>
 <b>4. Introduction.....</b>	 <b>13</b>
 <b>5. Review of the literature .....</b>	 <b>15</b>
5.1 Otorhinolaryngological services, audiology, and research in resource-poor context .....	15
5.2 Global burden of human rhino- and enteroviruses in respiratory infections .....	16
5.3 Chronic suppurative otitis media in children of developing countries .....	18
5.3.1 Etiology, epidemiology, and management of chronic otitis media .....	18
5.3.2 Course and consequences of chronic otitis media.....	20
5.4 Otorhinolaryngological signs and complications of sickle-cell disease .....	21
5.4.1 Epidemiology, pathophysiology, and management of sickle-cell disease .....	21
5.4.2 Hearing loss in sickle-cell disease.....	23
5.4.3 Other otorhinolaryngological manifestations in sickle-cell disease .....	24
5.5 Otorhinolaryngological pathology in human immunodeficiency virus –infected children .....	24
5.5.1 Pathophysiology, epidemiology, and management of human immunodeficiency virus infection .....	24
5.5.2 Otorhinolaryngological aspects of Human immunodeficiency virus infection.....	27

<b>6. Aims of the study .....</b>	<b>28</b>
<b>7. Children and methods.....</b>	<b>29</b>
7.1 Study groups and laboratory methods.....	29
7.2 Interview and physical examination.....	30
7.3 Otorhinolaryngological examination .....	30
7.4 Hearing measurements .....	32
7.5 Statistical analysis .....	34
<b>8. Results .....</b>	<b>35</b>
8.1 Respiratory infections – virology and clinical profiles [I].....	35
8.2 Chronic suppurative otitis media – clinical and microbiological variables [II].....	36
8.3 Sickle-cell disease-associated otorhinolaryngological pathology [III] .....	36
8.4 Human immunodeficiency virus-associated otorhinolaryngological pathology [IV] .....	39
<b>9. Discussion.....</b>	<b>40</b>
9.1 The role of picornaviruses in tropical respiratory infections .....	40
9.2 Middle-ear pathology .....	41
9.2.1 Otitis media as a complication of respiratory infection .....	41
9.2.2 Chronic suppurative otitis media.....	41
9.2.3 Middle-ear pathology in sickle-cell disease .....	42
9.2.4 The burden of otitis media in human immunodeficiency virus–positive children .....	43
9.3 Otorhinolaryngological profiles in different patient groups .....	44
9.4 Hearing Loss.....	45
9.4.1 Hearing loss due to chronic suppurative otitis media.....	45
9.4.2 Hearing loss in sickle-cell disease.....	45
9.4.3 Hearing problems in human immunodeficiency virus-positive children.....	46
9.5 Practical and ethical viewpoints.....	46

<b>10. Summary and conclusions .....</b>	<b>49</b>
<b>11. Acknowledgments .....</b>	<b>51</b>
<b>Kiitokset .....</b>	<b>53</b>
<b>12. References.....</b>	<b>55</b>
<b>Appendix .....</b>	<b>71</b>



# 1. ABSTRACT

**Background:** Otorhinolaryngological (ORL) diseases and hearing loss frequently occur among the children of developing countries, but they remain poorly characterized. A need therefore exists for more comprehensive understanding of the background and clinical features of these diseases in resource-poor settings.

**Patients and methods:** Voluntary paediatric outpatients of various specialties were examined at Hospital Pediátrico in Luanda, Angola. Study participants underwent medical history-taking, a thorough physical and ORL examination, hearing screening by brainstem auditory-evoked potentials (BAEP), and, at age 5 or older, pure-tone audiometry. Nasopharyngeal smears (NPS) were obtained from 102 children with respiratory symptoms and screened by polymerase chain reaction for human rhino- (HRV) and enterovirus (HEV). We took 18 bacterial culture smears from children with chronic suppurative otitis media (CSOM). Clinical data collected were compared between 23 children with CSOM vs. 23 age- and gender-matched controls, between 61 children with sickle-cell disease (SCD) vs. 61 controls, and between 78 human immunodeficiency virus (HIV) –positive children vs. 78 controls.

**Results:** In virus screening of 102 NPS specimens, 37 (36%) were positive: 27 (26%) for HRV alone, 3 (3%) for HEV alone, and 7 (7%) for HRV+HEV. HRV prevalence was highest during the coolest month, July, 47% (26/53), compared to 22% (8/49) ( $p=0.021$ ) in April-to-June. In the CSOM study, HIV positivity occurred in 14/22 (64%) of the CSOM children vs. none of the controls ( $p<0.001$ ) and tuberculosis in 8/22 (36%) vs. none ( $p=0.002$ ). The most frequent CSOM pathogens were *Proteus* (8/18, 44%) and *Pseudomonas* species (4/18, 22%). CSOM resulted in hearing loss of  $>25$  dB HL in pure-tone averages or BAEP in 17/33 (52%) of the affected ears. In the SCD study of 61 SCD children vs. 61 controls, bilateral hearing loss of  $>25$  dB HL at any frequency occurred in 9/25 (36%) SCD children vs. 3/28 (11%) controls ( $p=0.047$ ), whereas the prevalence of other ORL findings showed no significant difference. Of 78 HIV-positive children vs. 78 controls, ORL pathology was present in 72 (92%) HIV-positive vs. 61 (72%) control children ( $p=0.022$ ). Dental caries occurred in 44 (56%) of the HIV-positive children vs. 25 controls (32%) ( $p<0.001$ ), cervical lymphadenopathy  $>1$  cm in 35 (45%) vs. 8 (10%) ( $p<0.001$ ), facial skin lesions in 25 (32%) vs. 4 (5%) ( $p<0.001$ ), CSOM in 21 (26%) vs. 3 (4%) ( $p<0.001$ ), and bilateral hearing loss of  $>25$  dB HL in PTA or BAEP in 10 (13%) vs. one (1%) ( $p=0.009$ ).

**Conclusions:** In children of tropical Luanda, HRV and HEV emerged during respiratory symptoms. CSOM was associated with co-morbidity and hearing loss. Hearing loss occurred in SCD. ORL manifestations of HIV infection were common and various. In resource-poor settings, ORL and hearing services deserve a higher priority.



## YHTEENVETO

**Taustaa:** Korva-, nenä- ja kurkkutaudit (KNK) sekä kuulovauriot ovat kehitysmaiden lapsilla yleisiä mutta puutteellisesti tutkittuja. Siksi näiden sairauksien tausta ja kliininen kuva tarvitsevat tarkempaa määrittelyä.

**Potilaat ja menetelmät:** Tutkimukseen osallistui vapaaehtoisia eri erikoisalojen poliklinikkapotilaita Angolan pääkaupungin Luandan lastensairaala (Hospital Pediátrico). Osallistujilta kartoitettiin lääketieteelliset taustatiedot, tarkastettiin yleis- ja KNK-status sekä tutkittiin kuuloa aivorungon herätevasteiden ja vähintään 5-vuotiailta ilmajohteisen äänesaudiometrian avulla. Sadaltakahdelta hengitystieoireiselta lapselta otettiin nenänielusta tikkunäyte, josta tutkittiin polymeeraasiketjureaktiolla rino- ja enteroviruksia. Kahdeksaltatoista kroonista märkäistä välikorvatulehdusta (chronic suppurative otitis media, CSOM) sairastavalta lapselta otettiin bakteeriviljelynäyte. Tutkimustietoa kerättiin ja vertailtiin 23 CSOM:ta sairastavalta lapselta ja 23 ikä- ja sukupuolijakaumaltaan samankaltaiselta verrokilta; 61 sirppisoluanemiaa sairastavalta ja 61 verrokilta, sekä 78 HIV-infektiota sairastavalta ja 78 verrokilta.

**Tulokset:** Virusnäytteistä 37/102:ssa (36 %) todettiin virus: 27:ssä (26 %) pelkkä rinovirus, kolmessa (3 %) pelkkä enterovirus ja seitsemässä (7 %) molemmat virukset. Rinoviruksen esiintyvyys oli korkeimmillaan viimeisen heinäkuun aikana 47 % (26/53) ja muina kuukausina (huhti-kesäkuu yhdistettynä) 22 % (8/49) ( $p=0.021$ ). CSOM-tutkimuksessa HIV-infektio todettiin 14/22 (64%) CSOM-potilaalla eikä yhdelläkään kontrollilapsella ( $p<0.001$ ), ja tuberkuloosi 8/22:lla (36 %) CSOM-lapsella eikä yhdelläkään kontrollilla ( $p=0.002$ ). CSOM oli tyypillisesti *Proteus*- (8/18, 44 %) tai *Pseudomonas* -lajien (4/18, 22 %) aiheuttama, ja puhetaajuuksien 0.5-4 kHz kuulovaurio ( $>25$  dB HL) ilmeni 52 %:ssa vuotavista korvista. Sirppisoluanemiasta kärsivillä lapsilla molemminpuolinen, yleensä lievä kuulovaurio oli 9/25:llä (36 %) ja kontrollilapsista 3/28:lla (11 %) ( $p=0.047$ ); muuten ryhmien välillä ei KNK-löydösten esiintyvyyksissä ollut merkitseviä eroja. Poikkeavia KNK-löydöksiä oli 72/78:lla (92 %) HIV-positiivisella vs. 61/78:lla (72 %) kontrollilapsella ( $p=0.022$ ); kaulan alueen imusolmukesuurentumia 35 (45 %) HIV-positiivisella vs. kahdeksalla (10 %) kontrollilapsella ( $p<0.001$ ), iholöydöksiä 25:llä (32 %) vs. neljällä (5 %) ( $p<0.001$ ), CSOM 21:llä (26 %) vs. kolmella (4 %) ( $p<0.001$ ) ja molemminpuolista kuulonalemia ( $>25$  dB HL) 10:llä (13 %) vs. yhdellä (1 %) ( $p=0.009$ ).

**Johtopäätelmät:** Trooppisen Luandan lapsilla tavattiin rino- ja enteroviruksia hengitystieoireiden yhteydessä. Pitkittynyt märkävuoto välikorvasta liittyi usein

yleissairauksiin ja kuulonalenemaan. Sirppisoluanemiaa sairastavilla lapsilla esiintyi kuulo-ongelmia. HIV-positiivisista lapsista valtaosalla oli useita KNK-poikkeavuuksia. Kehitysmaiden lasten KNK- ja kuulo-ongelmiin olisi kiinnitettävä enemmän huomiota.

## 2. ORIGINAL PUBLICATIONS

This thesis is based on the following articles, which are referred to in the text by Roman numerals [I-IV].

- I. Taipale A, Pelkonen T, Roivainen M, Kaijalainen S, Bernardino L, Peltola H, Pitkäranta A. Human rhino- and enteroviruses in children with respiratory symptoms in Luanda, Angola. (Accepted by Paediatrics and International Child Health.)
- II. Taipale A, Pelkonen T, Taipale M, Bernardino L, Peltola H, Pitkäranta A. Chronic suppurative otitis media in children of Luanda, Angola. *Acta Paediatr* 2011; 100: 84-88.
- III. Taipale A, Pelkonen T, Bernardino L, Peltola H, Pitkäranta A. Hearing loss in Angolan children with sickle-cell disease. *Pediatr Int* 2012; 54: 854-857.
- IV. Taipale A, Pelkonen T, Taipale M, Roine I, Bernardino L, Peltola H, Pitkäranta A. Otorhinolaryngological findings and hearing in HIV-positive and HIV-negative children in a developing country. *Eur Arch Otorhinolaryngol* 2011; 268: 1527-1532.

Reprinted here (II–IV) with permission of the publishers.

### 3. ABBREVIATIONS

AOM	Acute otitis media
BAEP	Brainstem auditory-evoked potentials
CSOM	Chronic suppurative otitis media
HEV	Human enterovirus
HIV	Human immunodeficiency virus
HRV	Human rhinovirus
MEE	Middle-ear effusion
NPS	Nasopharyngeal smear
ORL	Otorhinolaryngological
PTA	Pure-tone average
PCR	Polymerase chain reaction
RTI	Respiratory tract infection
SCD	Sickle-cell disease
TM	Tympanic membrane
WHO	World Health Organization

## 4. INTRODUCTION

In developing countries, otorhinolaryngological (ORL) diseases and hearing loss are highly common. Despite their frequency, they remain, however, widely neglected. Some developing countries have no ORL specialists at all, and even the best served areas cannot be compared to the industrialized world. The shortage of ORL services includes not only medical specialists, but also the ORL knowledge of local general practitioners and nurses, as well as the facilities and technical equipment required. The situation is particularly poor in the audiology field. The brain-drain phenomenon, challenging the health care of many underserved regions, also extends to otolaryngology.<sup>1,2</sup>

This imbalance sets the young populations of resource-poor countries in an especially disadvantaged position, as many ORL diseases typically occur in childhood. Furthermore, the course of these diseases is often prolonged and complicated in children affected by poor hygiene, deficient nutrition, and chronic systemic illnesses. In 1996, the World Health Organization (WHO) held an expert meeting to develop strategies for prevention of deafness due to chronic otitis media in developing countries,<sup>3</sup> followed by a workshop concerning hearing aids a couple of years later.<sup>4</sup> Unfortunately, the medical and political awareness of the impact of untreated ORL diseases on the resource-poor societies has remained low.

Today, medical research concerning developing countries is still largely concentrated on life-threatening conditions such as severe infections. Meanwhile, data on basic epidemiology and the clinical picture of many common ORL diseases are sparse, noncomprehensive, or lacking. However, advances in health care have provided a better prognosis for huge numbers of underprivileged children, making research and management of their less fulminant diseases even more relevant.

The present study was launched in Luanda, the capital of Angola, to achieve epidemiological and clinical data on childhood ORL diseases, specifically related to picornavirus infections, CSOM, sickle-cell disease (SCD), and human immunodeficiency virus (HIV). The republic of Angola is located in southwestern Africa (Figure 1) and is currently experiencing a reconstruction era after its devastating 27-year civil war ending in 2002. Formerly part of the Portuguese Overseas Empire and a major sufferer from the Atlantic slave trade, the country became independent in 1975. Its steady economic growth, mainly fuelled by its rich diamond and petroleum resources, has, however, failed to benefit the vast majority of the population, such as the millions of inhabitants of the shantytowns of Luanda set up by inland escapees from the civil war (Table 1).<sup>5</sup>

In the present study, it very soon became evident that medical research in resource-poor settings has as typical challenges how to import the required equipment



Figure 1. Climate map of Africa, showing location of Luanda (figure by author).

Table 1. Comparison of country profiles of the Republic of Angola and Finland.<sup>5</sup>

	Angola	Finland
Population	19.1 million	5.3 million
Urbanized proportion of population	59%	85%
Life expectancy at birth	52 years	79 years
Under-five mortality rate	161/1000	3/1000
Number of HIV-positive children (estimated)	22 000	-
Conceptions affected by SCD (estimated)	16/1000	-
Gross national income per capita	3 960 US\$	47 170 US\$
General government expenditure on health	5.3%	11.7%
Proportion living under the international poverty line (1.25 US\$/day)	54%	Unknown
Adult literacy rate	70%	100%

through customs, how to recruit and communicate with people who are commonly unfamiliar both with the concept and purpose of research and with basic medical knowledge, how to manage problems of electricity supply and electronic devices without technical expertise available, and how to make short-term research locally useful and ethically sustainable. These challenges may in part explain the persistence of medical inequity between developed and developing world, but it should by no means prevent the worldwide efforts to reduce it – may the present study thus join in the mission with its own little share.

## 5. REVIEW OF THE LITERATURE

### 5.1 Otorhinolaryngological services, audiology, and research in resource-poor context

Even though the developing world stretches over virtually all the populated continents, the similar limitations in living conditions and health care systems in this vast area make the emerging ORL problems surprisingly similar. Therefore, in spite of the huge geographical, ethnic and cultural differences, it is no mere generalization to use as a common denominator “developing countries” for these regions. Typically, developing countries foster young, rapidly growing populations with broad ethnic and cultural diversity which suffer from high income inequalities. Domestic infrastructure, education, and manufacturing capacity seldom meet the demands of health care for these populations.<sup>6</sup> In practice, the restricted resources are often directed toward the most severe and lethal conditions, whereas diseases leading to disability but only rarely to immediate mortality are left with little attention.<sup>7</sup>

The developing world is therefore characterized by a marked shortage of ORL services, personnel, and training opportunities. For example, in a survey covering 18 African countries (not including Angola) with known public ORL services, the number of people per one ORL specialist ranged from 240 000 (South Africa) to 10 million (Malawi), in comparison to the 1:100 000 in the UK, which, in fact, ranks at the bottom among Western countries.<sup>8</sup>

Of these 18 African countries, 10 offered national ORL training, 2 had a training program for audiologists, and 12 had no or very poor ORL services outside major cities. The availability of various services was zero or poor as follows: audiology in 14 of the 18 participating countries, brainstem auditory-evoked potentials (BAEP) measurement in 16, hearing screening at schools in 15, hearing aids in 16, bone-anchored hearing aids in 17, cochlear implants in all 18, myringotomies/ventilation tubes in 14, tympanoplasty in 12, and mastoidectomy in 12. The authors concluded that in order to reach the lowest European level for ORL services, nearly 4800 additional ORL specialists are required in sub-Saharan Africa, and proposed establishing a worldwide “Developing World Forum for ENT Surgery, Audiology, and Speech Therapy” to reinforce the clinical, teaching, and research resources in ORL in developing countries.<sup>1,2</sup>

Annually, nearly 800 000 babies (6/1000 live births) are born in developing countries are now suffering or eventually will suffer from prelingual hearing loss.<sup>7,9</sup> By the age of nine, hearing loss prevalence will probably at least double.<sup>7</sup> Roughly

half these cases are believed to be preventable, and otitis media emerges as a major avoidable etiology alongside meningitis, cerebral malaria, measles, and mumps.<sup>7,9,10</sup> Usually, public awareness of the signs, consequences, and rehabilitation of childhood hearing loss remains low, and, in the absence of widespread hearing screening programs, the handicap may lead to permanent limitations in their linguistic and educational development, economically burdening already poor societies.<sup>3,7,10</sup> Outside the Western countries, only China, India, and Botswana produce hearing aids locally; generally, retail prices of imported devices remain unaffordable for customers in developing countries.<sup>11</sup>

Nationwide, systematic hearing screening programs are lacking in developing countries. Regional neonatal hearing screening programs have recently been launched in some resource-restricted areas,<sup>12</sup> but generally hearing-screening in developing countries relies on conventional audiometry typically performed in schools.<sup>9</sup> Simple low-cost, hand-held audiometers are available in developing-country settings.<sup>9,10</sup> In addition, screening takes place in immunization clinics.<sup>13</sup>

To save resources, screening has been targeted at known risk groups.<sup>12,14</sup>

Children at risk for hearing problems have also been identified by use of specific questionnaires on known risk factors and the guardian's observations of the child's speech development and response to sounds.<sup>15,16</sup> Increasingly, physiological measurements such as BAEP and otoacoustic emissions have also become available in resource-restricted countries.<sup>17</sup>

In general, relatively few ORL studies have been conducted in developing countries. In the absence of local databases of childhood ORL diseases, large population-based epidemiological surveys appear rarely and are available only from certain regions. Such reports typically concerned hearing loss and ear pathology in school-children. No cohort-based studies exist. Epidemiological data on ORL diseases in younger children also prove difficult to find. Since long-term follow-up was typically impossible, prevalence rather than incidence rates were generally reported for ear and hearing pathology. Another type of report available was hospital-based, comprising retrospective chart-reviews and prospective studies, mainly conducted in tertiary referral centres of major cities where local ORL services exist. These studies therefore probably concerned the more severe and complicated cases, or cases occurring in generally ill children, because in developing countries medical aid for ORL diseases is often sought only as a last resort.

## 5.2 Global burden of human rhino- and enteroviruses in respiratory infections

Human rhino- (HRV) and enteroviruses (HEV) are major pathogens in paediatric respiratory tract infections (RTI) in temperate climates of the northern hemisphere.<sup>18-23</sup> Being members of the *Enterovirus* genus and *Picornaviridae* family and thus



being only 30 nanometers in diameter, HRV and HEV carry single-stranded RNA inside an icosahedral, non-enveloped capsid consisting of four glycoproteins (VP1-4).<sup>24-26</sup> HRV is further classified into 3 (HRV-A, HRV-B, and HRV-C) and HEV into 4 (HEV-A-D) species, comprising more than 100 HRV and HEV (sero)types.<sup>25,27</sup>

The vast majority of HRV and several HEV types enter human cells through the intercellular adhesive molecule (ICAM-1) receptor<sup>25,28</sup> in the oral mucosa or upper respiratory epithelium. HRV is transmitted through contact with contaminated hands or with inanimate surfaces,<sup>20,22,29</sup> where it may persist for hours, even days.<sup>29,30</sup> HRV and HEV may be identified by cell culture or reverse transcription polymerase chain reaction (PCR), the latter being superior in sensitivity.<sup>20,31</sup>

The clinical profile of HRV infection involves the typical common cold symptoms of rhinorrhea, sneezing, and cough, and appear either alone or in combination with bacteria, and also rhinosinusitis, otitis media, and pneumonia.<sup>22,24,27,32-34</sup> In addition, HRV associates with asthma exacerbation in children.<sup>35,36</sup> HRV, being acid-sensitive, is unable to survive in the low gastric pH.<sup>20,22,27</sup> Some recent data suggest that HRV-C may enter the circulation,<sup>37</sup> but typical HRV infections are restricted to respiratory tissues. HEV also typically first invades the respiratory tissues, but, apart from common cold symptoms, it may equally likely cause gastroenteritis, “foot-and-mouth disease”, and severe, even fatal infections such as myocarditis and meningoencephalitis.<sup>24,38,39</sup>

In a child living in a temperate climate, picornaviruses cause an estimated five symptomatic infections per year,<sup>40</sup> and account for approximately half of the yearly common cold episodes.<sup>31,40</sup> Their incidence is, however, seasonal, with studies reporting proportions of 25 to 71% for HRV,<sup>18,19,23,32</sup> and 10 to 18%<sup>18,23</sup> for HEV in common colds. In temperate regions, HRV occurs throughout the year, but its incidence tends to peak in autumn,<sup>40-44</sup> and in some regions, also in spring.<sup>42-44</sup> Factors contributing to increased HRV transmission include high relative humidity<sup>45,46</sup> and crowding of children indoors during school terms.<sup>22,43</sup>

In general, little data exist on the seasonality of HEV-caused common colds. HEV serotypes responsible for more severe diseases may either emerge in an epidemical pattern, prevail in summer-autumn, or express no clear seasonality,<sup>24,38</sup> and generally no seasonality have been recorded for common colds from HEV.<sup>41,47</sup>

Data on common colds in tropical and subtropical climates are limited. However, the reported prevalences for HRV and HEV among children with respiratory symptoms range from 22 to 52%<sup>48-51</sup> and 3 to 15%<sup>49,50,52</sup> among outpatients/follow-up cohorts (Table 2), and 6.5 to 41%<sup>48,53-55</sup> and 4 to 20%<sup>53,54</sup> among hospitalized children. In the tropics, HRV occurrence reportedly increases with humidity or rainfall<sup>53,54</sup> and cooler temperatures,<sup>53</sup> and HEV with dryness and hot temperature,<sup>54</sup> but several studies found no clear seasonality pattern for HRV<sup>48,49,55</sup> or HEV.<sup>49,50,53,56</sup>

Picornaviruses are highly associated with otitis media in children: HRV has occurred in nasopharyngeal or middle ear samples of children with acute otitis media (AOM) in 20 to 41%<sup>18,41,57,58</sup> and HEV in 6 to 25%.<sup>41,58,59</sup> On the one hand, a follow-up

Table 2. Human rhinovirus (HEV) and human enterovirus (HEV) in paediatric outpatients of subtropical/tropical low-resource countries in the polymerase chain reaction era.

Country, year	N	Age (years)	Sampling method	HRV (%)	HEV (%)
Mozambique 2011 <sup>49</sup>	333	0–1	NPA	26	3
Kenya 2012 <sup>48</sup>	254	0–12	NPA	22	unmeasured
Kenya 2010 <sup>51</sup>	197	5–10	NPS	26.4	unmeasured
Senegal 2010 <sup>220</sup>	82	0–5	NPS	9.7	0

NPA, nasopharyngeal aspirate

NPS, nasopharyngeal swab

study of children under age two detected picornaviruses in MEE or nasopharyngeal samples of 54% of all children with AOM,<sup>41</sup> while on the other hand, another study showed that 30 to 34% of picornavirus colds were accompanied by AOM.<sup>18</sup> No data emerged on otitis media in picornavirus colds in children in tropical climates.

### 5.3 Chronic suppurative otitis media in children of developing countries

#### 5.3.1 ETIOLOGY, EPIDEMIOLOGY, AND MANAGEMENT OF CHRONIC OTITIS MEDIA

Chronic suppurative otitis media (CSOM) is a persistent inflammation of the mucosa and submucosa of the middle-ear cleft which produces a suppurative discharge through a chronically perforated tympanic membrane (TM).<sup>10,60–63</sup> The WHO defines otorrhoea as chronic after it endures for at least 2 weeks,<sup>10</sup> but no concerted opinion exists on the cut-off duration, and various values up to 3 months have been suggested.<sup>3,61,64</sup> The bacterial etiology of CSOM, reflecting either primary infection or secondary colonization, involves *Staph.aureus* and *P.aeruginosa*,<sup>10,60,65–69</sup> but children from a low-hygiene environment also exhibit enteric (e.g. *E.coli*) and low-virulence (e.g. *Proteus*) Gram-negative rods.<sup>60,70–73</sup> In addition, anaerobes may contribute in up to 71% of cases.

In the antibiotic era, CSOM has become a disease of the developing world. In addition, certain subpopulations in economically developed areas, such as the Inuits, South Pacific islanders, and Australian aboriginals, exhibit particularly high CSOM prevalences: up to at least 15%.<sup>74–78</sup> The vast majority, approximately 85 to 90%, of the global CSOM burden lies on Asia and South Pacific, mainly due to the large population of these areas, whereas the continent of Africa constitutes 4 to 8% of the cases worldwide. The WHO classifies CSOM prevalences as the following: the lowest <1%, low 1 to 2%, high 2 to 4%, and highest >4%; with prevalences exceeding 2% demanding urgent attention and >4% representing “a massive public health problem”.<sup>10</sup> Sub-Saharan Africa has reported prevalences of 1.1 to 4.6%,<sup>10,79–82</sup>




Figure 2. Risk factors, features, and events contributing to progression of otitis media towards chronic and complicated forms in children of developing countries.

contrasted with the European WHO estimate of 0.4%.<sup>10</sup> In Angola, in the 1980's, chronic otitis media occurred in 3.4% of schoolchildren.<sup>79</sup> Despite its relatively low global share in CSOM cases, of all deaths due to otitis media complications in 2008, Africa accounted for two-thirds.<sup>83</sup>

Approximately one-fifth of the CSOM cases occur in children less than age 5.<sup>84</sup> Risk factors for chronic otorrhea include poor socio-economic status, crowding, malnutrition, short duration of breastfeeding, parental smoking, and lack of clean water for washing.<sup>72,75,77,78,82,85</sup> In resource-poor settings, deficiencies in vitamin A and zinc may also add to the risk for a chronic or more severe course of otitis media.<sup>86-88</sup> In addition to the living conditions and environmental variables, genetic factors predispose to chronic otitis through anatomical (e.g. short auditory tube)<sup>60,63</sup> and functional (e.g. epithelial ciliary motility) variance<sup>60</sup> (Figure 2).

In general, few data exist on treatment of paediatric CSOM in developing countries. The strongest evidence concerns the resolution of otorrhea, while in the lack of long-term follow-up, data on recurrence, restoration of the tympanic membrane and hearing, complications, and adverse effects of the treatment remain uncertain.<sup>89</sup> However, the safety of topical ofloxacin is evident in children.<sup>90</sup>

In resource-poor settings, the treatment is based on regular ear cleansing, which in itself has no therapeutic efficacy,<sup>91</sup> but facilitates the entry of topical antimicrobials into the middle ear.<sup>92</sup> Clinical trials have shown the efficacy of ototopical an-

tibiotics in treating CSOM in children;<sup>89</sup> at least ofloxacin,<sup>70,91,93</sup> ciprofloxacin,<sup>68,94</sup> and neomycin-polymyxin-B<sup>93</sup> have been studied in low-resource settings. Other topical antibiotics in clinical use include gentamycin, tobramycin, chloramphenicol, trimethoprim, and erythromycin.<sup>10</sup> Adding systemic antibiotics to local treatment is not recommended due to its lack of efficacy and low cost-effectiveness.<sup>10</sup> Topical antiseptics, such as boric acid, are cheap and useful,<sup>95</sup> but remain less efficacious and may cause more local irritation than a local antibiotic would.<sup>94</sup>

### 5.3.2 COURSE AND CONSEQUENCES OF CHRONIC OTITIS MEDIA

Generally, CSOM is regarded as forming a continuum with AOM and otitis media with effusion,<sup>60,61,63</sup> and in developing countries it is mostly believed to arise from an unresolved or untreated AOM episode, whereas traumatic TM perforations may play a lesser role.<sup>3,10,63,96</sup> In industrialized countries, CSOM usually associates with tympanostomy tubes.<sup>97,99,98,99</sup> In otitis-prone children, early bacterial colonization, leading to formation of biofilms and chronic inflammation in the nasopharynx and auditory tube, may contribute to the persistence of middle-ear infections.<sup>100,101</sup>

Prolonged inflammation of the middle-ear cavity induces changes in the surrounding structures that complicate treatment attempts and the healing process. In the mucosal lining, metaplasia occurs, and mucus-producing glandules, as well as ciliated cells and highly vascular granulation tissue emerge. Ongoing infection may also invade the underlying bone, causing osteitis and reactive osteogenesis. Pathological changes detectable in TM include fibrosis, thickening, retraction, epithelialization, and myringitis that may lead to polyp formation. Cholesteatoma may also occur.<sup>10,61,62,102,103</sup>

CSOM may lead to conductive hearing loss resulting from the discharge in the ear canal. Thickening, retraction, and perforation of the eardrum also produce conductive hearing loss.<sup>3,61</sup> In addition, resorption and disruption of the ossicular chain, in 11 to 68% of patients of various ages discovered during surgical treatment,<sup>104-106</sup> may worsen hearing loss. Bacterial toxins may also enter the inner ear through the round window membrane, causing cochlear damage and a sensorineural or mixed type of hearing loss.<sup>107-109</sup> Another risk factor for cochlear hearing loss is ototoxicity from topical aminoglycosides, which are usually preferred to the safer but more expensive quinolones in lower-resource health care.<sup>3</sup> However, ototoxicity may not be very usual in clinical settings.<sup>110</sup>

Apart from hearing loss, CSOM may lead to suppurative complications that can be extracranial, such as labyrinthitis, mastoiditis/mastoid abscess, petrositis or facial paralysis; or intracranial, such as thrombosis of lateral or cavernous sinus, meningitis, cerebellitis, epidural/subdural/cerebral abscess, hydrocephalus, and labyrinth sclerosis. These intracranial complications hold the main responsibility

for CSOM-associated mortality, which in Africa has been estimated by the WHO to reach 1.8%.<sup>10,84,102,103,111,112</sup>

In sub-Saharan Africa, up to 3100 children under 15 thus far have died annually in the 2000's due to complications of otitis media.<sup>84</sup> Even though severe complications of CSOM have been estimated to be relatively rare in children,<sup>10</sup> their risk is significantly higher than for adults.<sup>84,111</sup> Among CSOM patients presenting at ORL clinics in Africa, subperiosteal abscesses and intracranial complications have been reported in 6 to 11%.<sup>103,111,112</sup> Apart from mortality, complications of CSOM may lead to severe permanent neurological sequelae.<sup>10</sup>

In Thailand, 102 (0.59%) complications occurred in 17 144 otitis media cases seen in health care settings in the 1980's, and the majority of them arose from CSOM. In that series, the most frequent extracranial manifestations were facial paralysis, subperiosteal abscess, and labyrinthitis, whereas the intracranial complications were mainly meningitis and intracranial abscess, leading to death in 18.6%.<sup>113</sup>

However, chronic suppuration may also spontaneously resolve, followed by healing of the perforation or by chronic perforation with either no or only an occasional discharge.<sup>63</sup> In follow-up of CSOM children without therapy, suppuration had ceased at 6 weeks in 18%<sup>72</sup> and at 16 weeks in 22%<sup>91</sup>; and only 13% had intact TMs at 16 weeks.<sup>91</sup>

Of the worldwide otitis media-related hearing loss, 90% occurs in developing countries, where CSOM appears a major cause for acquired hearing loss in children.<sup>3,80,96,114</sup> Among paediatric CSOM patients, hearing loss has been documented in 47 to 96%,<sup>80-82,114-116</sup> and mainly presents as slight or moderate.<sup>68,81,82,114</sup> CSOM affects most heavily young children,<sup>3,72,85,112,115</sup> and may endanger their language acquisition, social development, and scholastic performance.<sup>75,82,117</sup>

## **5.4 Otorhinolaryngological signs and complications of sickle-cell disease**

### **5.4.1 EPIDEMIOLOGY, PATHOPHYSIOLOGY, AND MANAGEMENT OF SICKLE-CELL DISEASE**

Sickle-cell disease (SCD) is the most prevalent monogenic disease worldwide, with 300 000 affected children being born every year.<sup>118</sup> Of the patients, the developing countries have at least 80%.<sup>119</sup> In Sub-Saharan Africa, 180 000 births occur annually, and due to improved patient survival, the burden of disease in this area is increasing.<sup>119</sup> Given survival rates half the local average, six million SCD patients will live in the area.<sup>120</sup> The generally healthy carriers of the sickle-cell mutation, "trait", appear to be nearly 90% protected against severe *Plasmodium falciparum* malaria,<sup>121</sup> and the malaria-endemic regions of Arabia, Mediterranean, India, and Africa thus experience the highest SCD incidence.<sup>120,122-124</sup>




Figure 3. Abnormally shaped erythrocytes in sickle-cell disease (figure provided by specialist in clinical chemistry Päivi Helminen-Pacius from HUSLAB laboratory, Helsinki University Central Hospital, Finland).

In Africa, the top SCD rates occur between 15° north and 20° south latitudes where up to 40% of the population may carry the sickle-cell trait.<sup>125</sup> In Angola, SCD affects 1.6% of conceptions,<sup>123</sup> which may exceed the actual proportion of newborns, because SCD adversely affects fetal survival.<sup>126</sup>

SCD is a haemoglobin disorder resulting from coinheritance of the “sickle” allele, HbS, and another mutant haemoglobin allele; either a second HbS resulting in the homozygous HbSS disease called sickle-cell anemia, or another type of a mutant allele, such as HbC or thalassemia, resulting in compound heterozygous inheritance.<sup>120,124,127</sup> In sub-Saharan Africa, the HbSS genotype prevails, with only small minorities of HbSC and other types.<sup>122,125</sup>

The sickle haemoglobin contains a single amino acid substitution (valine for glutamic acid) in its  $\beta$ -chains, leading to loss of form and function of the red blood cells (Figure 3), chronic haemolytic anemia, and acute vaso-occlusive crises, as well as various systemic outcomes related to haemolysis and tissue infarctation.<sup>124,127</sup> Through vaso-occlusive congestion, repeated splenic infarcts, and fibrosis, SCD leads to functional hyposplenism,<sup>128</sup> which makes the children highly susceptible to severe bacterial infections, particularly by *Streptococcus pneumoniae*,<sup>123,129-132</sup> which markedly contributes to the estimated 50 to 90% mortality among affected African children.<sup>122,133,134</sup> In addition to the genetic factors, environment plays a marked role in the phenotype of SCD patients,<sup>135</sup> and a paucity of data exists on the complications and natural course of SCD in developing countries.<sup>119</sup>



Table 3. Hearing loss exceeding 25 dB HL in audiometry, and related middle-ear pathology (MEP) in children with sickle-cell disease. Adapted from: Taipale A, Pelkonen T, Bernardino L, Peltola H, Pitkäranta A. Hearing loss in children with sickle-cell disease. *Pediatrics International* 2012; 854-857. © The Authors. Pediatrics International © 2012 Japan Pediatric Society.

Author, country	N	Age (years)	Hearing loss		Proportion of MEP in hearing loss
			n (%)	bilateral	
Alabi et al., 2008 (139) Nigeria	80	4-15	25 (31%)	22	28% <sup>a</sup>
Odetoyinbo and Adekile, 1987 (144) Nigeria	56	6-15	12 (21%)	7	0%
Mgbor and Emodi, 2004 (143) Nigeria	52	6-19	7 (13%)	7	no mention
De Castro Silva et al., 2010 (140) Brazil	40	8-20	11 (28%) <sup>a</sup>	5	0%
Tsibulevskaya et al., 1996 (138) Kenya	33	7-14	12 (36%)	2	0%
MacDonald et al., 1999 (149) USA	84	8 months - 24 yrs	26% <sup>c</sup>	NM	23% <sup>a</sup>
Forman-Franco et al., 1982 (141) USA	54	12 (mean)	11%	NM	7%

<sup>a</sup> Middle-ear effusion

<sup>b</sup> Hearing loss exceeding 20 dB, measured by brainstem auditory-evoked potentials

<sup>c</sup> Neither number of children with nor definition of hearing loss mentioned

Successful management of SCD is possible in resource-poor settings and includes a balanced diet with folic acid supplement, avoidance of extreme physical stress, heat, and dehydration, treatment with penicillin prophylaxis, pneumococcal vaccinations, analgetics, hydroxyurea, and blood transfusions.<sup>118,125,136</sup> Newborn screening, not yet usual in developing countries, is necessary for early identification and management of new SCD children,<sup>137</sup> and has the potential to reduce SCD mortality.<sup>138</sup>

#### 5.4.2 HEARING LOSS IN SICKLE-CELL DISEASE

SCD is known to associate with sensorineural hearing loss in children,<sup>139-144</sup> adolescents,<sup>140,141,143-149</sup> and adults.<sup>146-152</sup> Hearing loss has been proposed to relate to cochlear ischaemia, based on post-mortem observations of the temporal bones of an SCD child with known hearing loss.<sup>153</sup> Corroborating this theory are case reports of adolescents with reversible hearing loss in association with vaso-occlusive crisis.<sup>141,154</sup> Other support for hypoxia in the pathogenesis of SCD-related hearing loss include the high dependency of cochlear homeostasis on its vascular supply, inner-ear vascularization deriving exclusively from the internal auditory artery, and the degeneration and ossification in the cochlea after internal auditory artery occlusion.<sup>155</sup> In one SCD child, labyrinthitis ossificans has been a cause of progressive sensorineural hearing loss.<sup>156</sup>

Hearing loss of varying definitions has occurred in SCD children/adolescents at a rate of 3.5 to 12% in developed countries.<sup>142,157-159</sup> In developing countries, conductive or sensorineural bilateral hearing loss exceeding 25 dB HL has been reported in 6 to 27.3% of mainly school-aged children (Table 3). Some studies have perceived higher prevalences of hearing loss in older patients.<sup>147,148,150,152,160</sup> The SCD-related hearing loss has been reported to involve either exclusively high,<sup>145,147-149,160</sup> or all<sup>142-144,146,154,160,161</sup> frequencies, with both sudden<sup>141,154</sup> and slow<sup>145,147</sup> onset. The sudden-onset hearing loss may relate to an acute vaso-occlusive event in the cochlea,<sup>141,154</sup> whereas the slow-onset type may result from anoxia of the hair cells produced by non-acute erythrocyte sickling and chronically impaired local oxygen balance.<sup>162</sup>

### 5.4.3 OTHER OTORHINOLARYNGOLOGICAL MANIFESTATIONS IN SICKLE-CELL DISEASE

The ORL manifestations of SCD remain poorly described in children. The estimation is that unvaccinated SCD children without prophylactic penicillin may experience a 33- to 100-fold<sup>123,132,163</sup> risk for invasive pneumococcal infections. *Streptococcus pneumoniae* frequently causes otitis media, which is, in contrast, usually a localized infection, and no studies have been aimed at defining the incidence or prevalence of otitis media in SCD children. However, antibody-deficient children without haemoglobin disorders do suffer from recurrent and persistent ear infections.<sup>164,165</sup>

SCD-related ORL findings, mainly described in case reports, include bone infarcts, osteonecrosis, osteomyelitis, extramedullary hematopoiesis in the paranasal sinuses, bone marrow hyperplasia of the maxilla and skull base, soft-tissue abscesses, and cervical lymphadenopathy.<sup>166-169</sup> Adenoid hypertrophy<sup>170,171</sup> and nasal obstruction resulting from turbinate congestion<sup>170</sup> may also occur in children. In the absence of prospective follow-up studies, the frequencies of various manifestations remain, however, unknown.

## 5.5 Otorhinolaryngological pathology in human immunodeficiency virus –infected children

### 5.5.1 PATHOPHYSIOLOGY, EPIDEMIOLOGY, AND MANAGEMENT OF HUMAN IMMUNODEFICIENCY VIRUS INFECTION

Human immunodeficiency virus (HIV) is an enveloped, single-stranded RNA retrovirus in the *Lentivirus* genus of the Retroviridae family. With the aid of its external gp120 glycoprotein, HIV enters CD4 (cluster of differentiation 4) glycoprotein-expressing human cells, e.g. helper T cells, monocytes, macrophages, and glia cells. Initial infection occurs through percutaneous or mucous membrane contact with contaminated objects or body fluids, as well as transplacentally in utero. Untreated



Table 4. World Health Organization clinical stages of human immunodeficiency virus (HIV) infection.<sup>177</sup>

Clinical stage	General findings	Otorhinolaryngological findings
I – Asymptomatic	<ul style="list-style-type: none"> <li>• Persistent generalized lymphadenopathy</li> </ul>	<ul style="list-style-type: none"> <li>• Persistent cervical lymphadenopathy</li> </ul>
II – Mild	<ul style="list-style-type: none"> <li>• Herpes zoster</li> <li>• Papular or seborrhoeic dermatitis</li> <li>• Extensive wart virus or molluscum contagiosum infection</li> <li>• Persistent hepatosplenomegaly</li> </ul>	<ul style="list-style-type: none"> <li>• Recurrent sinusitis, tonsillitis, otitis media, pharyngitis</li> <li>• Angular cheilitis</li> <li>• Lineal gingival erythema</li> <li>• Recurrent oral ulceration</li> <li>• Persistent parotid enlargement</li> </ul>
III – Advanced	<ul style="list-style-type: none"> <li>• Moderate malnutrition or wasting</li> <li>• Persistent fever or diarrhoea</li> <li>• Pulmonary or lymphatic tuberculosis</li> <li>• Recurrent bacterial pneumonia</li> <li>• Chronic HIV-associated lung disease</li> <li>• Anaemia, neutropaenia, or chronic thrombocytopaenia</li> </ul>	<ul style="list-style-type: none"> <li>• Acute necrotizing stomatitis or gingivitis</li> <li>• Persistent oral candidiasis</li> <li>• Oral hairy leukoplakia</li> <li>• Cervical lymph node tuberculosis</li> </ul>
IV – Severe	<ul style="list-style-type: none"> <li>• Severe malnutrition, stunting or wasting</li> <li>• Chronic cryptosporidiosis, isosporiasis, or herpes simplex infection</li> <li>• Recurrent severe bacterial infection</li> <li>• <i>Pneumocystis jirovecii</i> pneumonia</li> <li>• Atypical mycobacteriosis, extrapulmonary tuberculosis or cryptococcosis, disseminated mycosis</li> <li>• HIV encephalopathy</li> <li>• Kaposi sarcoma, Non-Hodgkin lymphoma</li> </ul>	<ul style="list-style-type: none"> <li>• Chronic herpes simplex infection (orolabial)</li> <li>• Oesophageal candidiasis</li> <li>• Kaposi sarcoma (oropharyngeal, intraparotid)</li> <li>• Non-Hodgkin lymphoma (intraoral, cervical)</li> </ul>

infection leads to deterioration of the host's immune system, subsequent severe bacterial and viral infections, diseases by opportunistic microorganisms (candida, *Pneumocystis jirovecii*, *Cryptococcus*, mycobacteria), occurrence of malignancies, general wasting, and, finally, death.<sup>172</sup>

In resource-poor countries where children are usually infected intrauterinally, perinatally, or via breastfeeding, roughly half the untreated children progress to death within the first 2 years of life,<sup>173,174</sup> while the rest appear to be slow progressors with a median survival of 6 years, or long-term survivors with several asymptomatic years.<sup>172</sup> Paediatric HIV is mainly being caused by the viral subtype HIV-1, whereas HIV-2 is of West African origin and occurs rarely in children.<sup>175</sup>

Worldwide, sub-Saharan Africa carries the heaviest burden of HIV. In 2009, 2.3 million HIV-positive children lived in the area, constituting more than 90% of

Table 5. Percentages of otorhinolaryngological findings in representative studies in HIV-positive children. Adapted from: Springer and *European Archives of Oto-Rhino-Laryngology*, volume 268 (2011), p.1527-1532. Otorhinolaryngological findings in Human immunodeficiency virus-positive and -negative children in a developing country. Taipale A, Pelkonen T, Taipale M, Roine I, Bernardino L, Peltola H, Pitkäranta A., Table 4; with kind permission from Springer Science and Business Media.

Author, country	N	Cervical lymphadenopathy	Eczema	Chronic OM	Parotid hypertrophy	Oral candidiasis	Any ORL <sup>b</sup> pathology
Chaloryoo et al. (1998) <sup>186</sup> Thailand	250	42	23	18	5	60	NM <sup>c</sup>
Gondim et al. (2000) <sup>188</sup> Brazil	20 <sup>a</sup>	60	NM	15	25	35	90
Hadfield et al. (1996) <sup>185</sup> UK	66	71	15	8	14	33	91
Singh et al. (2003) <sup>184</sup> UK	107	36	NM	NM	6	18	50

OM, otitis media

ORL, otorhinolaryngological

NM, not mentioned

<sup>a</sup> Mean age 3.8 years

all paediatric HIV cases. Both the incidence of and mortality from paediatric HIV declined slightly in the area during the 2000's, resulting in a net increase in the number of children living with HIV. In fact, two-thirds of the world's HIV patients live in the area of 10 sub-Saharan countries including Angola. The number of Angolan HIV-positive children reached 22 000 in 2009, at a time when only some 15% of these children received antiretroviral therapy.<sup>176</sup>

In children at least 18 months old, HIV may be diagnosed serologically, whereas – due to the persistence of maternal antibodies a definite diagnosis in younger children requires virological tests.<sup>177</sup> A positive serological test in an infant, however, reveals exposure to HIV.<sup>178</sup> In low-resource health care, where the availability of virological testing is constrained, clinical signs<sup>178</sup> or other parameters<sup>179</sup> may be necessary to predict the HIV status of the infant. In developed countries, HIV staging is based on viral PCR and CD4 T cell counts, which are however not always available in resource-poor settings. Therefore, based on clinical signs, WHO has staged the severity of HIV infection in four categories (Table 4).<sup>177</sup>

Medical management of HIV infection is based on combination therapy with antiretroviral drugs, called, by their mechanism of action, nucleoside and non-nucleoside reverse-transcriptase inhibitors, and protease inhibitors. In children under age 2, antiretrovirals should be started promptly after diagnosis, whereas in older children, the beginning of treatment should be guided by CD4 counts.<sup>178</sup> Zidovudine prophylaxis in the newborn has been used for prevention of intrapar-

tum HIV transmission, but a two- or three-drug regimen has proven more efficacious if the mother has not received antepartum antiretroviral therapy.<sup>180</sup> However, in low-income settings, limitations in infrastructure, drug availability, monitoring resources, and treatment adherence challenge HIV prevention and management.<sup>175</sup>

### 5.5.2 OTORHINOLARYNGOLOGICAL ASPECTS OF HUMAN IMMUNODEFICIENCY VIRUS INFECTION

Worldwide, more than 90% of newly infected HIV children live in sub-Saharan Africa,<sup>176</sup> and at least 22 000 HIV-infected children live in Angola.<sup>181</sup> Paediatric HIV manifests in ORL area in multiple ways<sup>182-189</sup> and in up to 91% of the infected;<sup>184,185,188</sup> often already during the first years of life.<sup>188</sup> Table 5 summarizes previously documented frequencies of common ORL findings among HIV-positive children of developing countries, and Table 4 presents the ORL aspects of the HIV staging system. In addition to the manifestations mentioned in these tables, marked adenotonsillar enlargement may occur in HIV children,<sup>182</sup> as well as initial hypertrophy followed by end-stage atrophy.<sup>190</sup>

Hearing loss has been reported in HIV-positive children in 27.5 to 38.8%,<sup>191-194</sup> and sensorineural hearing loss in 7.1 to 16.7%.<sup>192-194</sup> Possible causes for sensorineural hearing loss in HIV patients include ototoxicity from antiretroviral therapy (mainly nucleoside analogues, as reported in adults), antituberculous medication (streptomycin),<sup>195,196</sup> aminoglycosides,<sup>195,197,198</sup> or possibly the direct neurotoxic effects of the HIV infection itself.<sup>192-194</sup> In addition, cytomegalovirus and *Cryptococcus* infections have induced hearing loss in HIV-positive adults,<sup>199,200</sup> with no reports in children.

Despite the possible presence of cochlear and neural lesions from these causes, however, a major contributor in HIV-positive children's hearing problems is otitis media, which strongly associates with HIV both in developed<sup>187,201-203</sup> and in developing<sup>191,204-206</sup> countries. The severity of immunodeficiency may correlate with a tendency for chronic<sup>204,207</sup> or recurrent<sup>202,203</sup> otitis media. In African HIV-infected children seen at hospital, 4 to 20% have presented with otorrhea,<sup>206,208,209</sup> which usually involves bacteria typical of CSOM in developing countries.<sup>186,210</sup> HIV-infected children are also subject to complications of otitis media.<sup>187</sup>

## 6. AIMS OF THE STUDY

The primary aim of the present study was to collect epidemiological and clinical data on paediatric ORL findings in Luanda, Angola. The specific aims were to assess:

- the prevalence of and clinical findings concerning human rhino- and enterovirus positivity during the cool season;
- CSOM-related background characteristics, bacteriology, and hearing;
- ORL status and hearing in SCD patients;
- HIV-associated ORL findings and hearing.

## 7. CHILDREN AND METHODS

### 7.1 Study groups and laboratory methods

The study data were collected in the Paediatric Hospital of Luanda (Hospital Pediátrico David Bernardino), a university teaching hospital, in January-July 2008. The Paediatric Hospital's ethics committee gave approval for the study on December 27<sup>th</sup> 2007. The study children were recruited among voluntary outpatients of the Paediatric Hospital's different polyclinics (pulmonology, surgery, general pediatrics, neurology, HIV, sickle-cell disease, gastroenterology, infectious diseases). In addition, a few accompanying siblings of the patients were included. At the general enrollment desk for all clinic outpatients, the receiving secretary provided basic information on the study and offered the opportunity for everybody to participate. Children with ear discharge, as well as outpatients of the HIV and SCD clinics were addressed in particular. In addition, a local nurse who served as a study assistant helped in the recruitment. All volunteering children were primarily accepted, and the children or their guardians then received more specific oral information on the aims and procedures of the study from the examining doctor prior to giving their oral consent.

During the study period, data came from a total of 406 children. All children meeting the inclusion criteria for Studies I to IV were then extracted from this data pool. In Studies II to IV, the age- and gender-matched controls for the patient groups were randomly selected among the remaining data pool. The control groups in Studies III and IV therefore included 29 children in common.

For Study I, 102 children displaying symptoms of the common cold (rhinorrhoea or cough, with or without fever) were recruited during April-July. Of these children, 79 were participants from Studies II to IV: 65 patients from Studies II to IV and 14 controls from Studies III and IV.

In Study I, a nasopharyngeal smear (NPS) was obtained with a sterile cotton swab from each of the 102 participants. The NPS specimens were preserved frozen, transported to Helsinki, Finland, and analyzed for HRV and HEV by real-time PCR assay. The analysis for HRV was performed as described earlier,<sup>211</sup> applying the cut-off threshold of 8% of the strongest positive control. The analysis for HEV was performed according to the method by the US Centers for Disease Control and Prevention, Atlanta, Georgia, USA<sup>212</sup> applying the cut-off threshold of 10% of the strongest positive control. The real-time-PCR analyses were executed at Finland's National Institute of Health and Welfare.

Study II comprised 23 children with CSOM and 23 age- and gender-matched controls without ear complaints or TM perforations. The CSOM group comprised

children with purulent middle-ear discharge for at least 2 months. Bacterial smears from the discharging ears – or in bilateral cases, from the more profusely discharging ears – were taken with a sterile cotton swab and cultured under aerobic conditions according to the norms of the National Committee for Clinical Laboratory Standards.

For Study III, 61 children with SCD from the SCD outpatient clinic were recruited; 61 age- and gender-matched children without clinical suspicion of SCD, known sickle cell trait, or a previous cerebral infection served as controls. SCD diagnosis was based on haemoglobin electrophoresis. The controls did not routinely undergo haemoglobin analysis.

Study IV comprised 78 HIV-positive outpatients from the HIV polyclinic, and an age- and gender-matched group of 78 controls without clinical suspicion of HIV. HIV was diagnosed with a specific antibody-detection test (Hexagon HIV® Human, Wiesbaden, Germany). The HIV negativity of the controls was not serologically verified.

## 7.2 Interview and physical examination

All study participants, together with their guardians, underwent a structured interview concerning the child's living conditions and family size, medical history and vaccinations, intake of medication, and ear- and hearing complaints. Subsequently, a basic physical examination and a thorough ORL status were performed (Appendix). The children's linguistic development was assessed in a free conversation with the child; especially with the youngest children, assessment was based on information from the guardian.

The basic physical examination comprised an overall assessment of the child's general condition, measurement of weight and height, cardiac and pulmonary auscultation, and palpation of the abdomen. In addition, the inguinal and axillar lymph nodes were palpated and dermal abnormalities registered. The children's nutritional status was assessed by the WHO standards for weight for age, and children with a low weight for age (below -2 SD or the 3<sup>rd</sup> percentile) were considered malnourished. Pneumonia was diagnosed in children with fever, cough, and tachypnea, augmented by pulmonary auscultation and, when available, chest x-rays.<sup>213,214</sup> The guardian or the examining doctor assessed the child's feverishness without a thermometer. The HIV-positive children who presented with clinical signs of WHO disease-stage III or IV<sup>177</sup> (Table 4) were considered as having an advanced state of the disease.

## 7.3 Otorhinolaryngological examination

All study children underwent a thorough ORL examination including anterior rhinoscopy, oral and oropharyngeal cavity examination, pneumatic otoscopy, and


tympanometry. Any facial abnormalities were visually assessed, and the size and appearance of auricles, as well as possible existence of preauricular fistules, were recorded. In addition, the cervical area was palpated, and the size (cm) and location of palpable lymph nodes or tumors were recorded, as well as the size and consistency of the salivary glands.

The oral and oropharyngeal examination included the oral mucosa, tonsils, and the dental status. Absence of teeth was noted, as well as abnormal development, and presence of cariotic lesions in the dentition. Tonsillar size was graded on the basis of Mallampati classification 0-4 (0: tonsils fit within the tonsillar fossa; 1: tonsils 0 to 25% of the space between the pillars; 2: tonsils 25 to 50% of the space between the pillars; 3: tonsils 50 to 75% of the space between the pillars; 4: tonsils 75 to 100% of the space between the pillars).

Otologic assessment comprised pneumatic otoscopy and tympanometry. During otoscopy, the ear canal was cleansed from cerumen or suppuration with cotton swabs and alligator forceps. However, if a satisfactory view of the tympanic membrane was impossible due to tightly impacted cerumen or excessive suppuration, the case was simply defined as “cerumen impaction” or “suppuration.” Diagnosis of AOM was based on detection of middle-ear fluid behind an inflamed TM, accompanied by any sign of local or general infection. Signs of fluid in the middle ear comprised visible fluid or pus behind the TM, flatness or bulging of the TM, restricted mobility of the TM. Inflamed TMs were red, yellow, or opaque. Local and general signs of infection included earache, fever, rhinorrhea, cough, and vomiting. Children with fluid in the middle ear but without inflamed TMs and evidence of an acute infection were defined as having middle-ear effusion (MEE). Suppurative discharge from the middle ear enduring for less than 2 months was also defined as AOM. Suppuration persisting for over 2 months was defined as CSOM.

Tympanometry was performed with a GSI 38 Auto Tymp tympanometer (Grason-Stadler, Inc., Eden Prairie, MN, USA). A pressure range of -400 to +200 daPa was tested with a 226-Hz probe. The tympanograms obtained were classified as A, B, and C according to the presence and location of the peak on the horizontal axis (daPa). (Figures 4 a-c) Curve gradients were not accounted for. Curves with a distinct peak at  $\pm 150$  to  $\pm 100$  daPa were designated as type A, which suggested normal pressure in the middle ear. Curves with a peak occurring at  $\pm 400$  to  $\pm 150$  daPa were designated as type C, and suggested possible negative pressure in the middle ear. Both types A and C were interpreted as indicating an intact TM without fluid in the middle ear. Peakless curves were designated as type B and considered to suggest either fluid in the middle ear, TM perforation, or cerumen impaction.

The final assessment of middle-ear status was based on both tympanometry and otoscopy. However, even if nonremovable cerumen prevented a view of the TM with an otoscope, obtaining a type A or C tympanogram was considered as ruling out TM perforation.



Figures 4a-c. Examples of tympanogram curve types A-C. ECV – external canal volume, normally in children approximately 0.6 cm<sup>3</sup>; GR – tympanogram gradient, width of graph at 50% of the maximal admittance; PEAK – the maximal (static) admittance of tympanic membrane.

a: Type A – graph with a distinct peak occurring at a pressure higher than -150 daPa (on x axis).

b: type B – flat graph with no detectable peak.

c: type C – graph with a distinct peak at a pressure less than -150 daPa.

## 7.4 Hearing measurements

Hearing measurements comprised BAEP threshold screening (Bera MADSEN Octavus® system v.2.001, Windows XP/2000 compatible; GN Otometrics, Taastrup, Denmark), and air-conduction pure-tone audiometry (Midimate 622 Clinical/Diagnostic Audiometer; Madsen Electronics, Taastrup, Denmark). Click-stimulated BAEP screening was applied in all children. Pure-tone audiometry was measured without a masking tone. Audiometry was performed in children of at least 5 years of age, due to its cognitive and co-operative requirements.

In pure-tone audiometry, hearing was tested at 0.125, 0.25, 0.5, 1, 2, 4, 6, and 8 kHz (Figure 5). Starting from 1 kHz, the higher frequencies were tested first in ascending order, followed by the lower frequencies in descending order. At each frequency, the screening level was 20 dB HL. If the child failed to respond at 20 dB HL, the sound level was increased by 5 dB until a response was obtained. If the result at any frequency was assessed as unreliable, the hearing threshold was confirmed by immediate re-testing. Pure-tone averages (PTA), the mean hearing thresholds of 0.5, 1, 2, and 4 kHz, were applied in Studies II and IV, whereas in Study IV each frequency in the range 0.25 to 8 kHz was analyzed separately. The hearing thresholds were graded according to the WHO scale (Table 6).<sup>215</sup> While performing pure-tone audiometry, the researchers continuously assessed the surrounding noise level. In the absence of sound-proof facilities, all sounds from inside the examination room, such as air-conditioning, were quieted, and during any noise from outside the room, the screening was paused.

BAEP thresholds were measured at 40, 60, and 80 dB, (Figure 6) and the results interpreted as follows: absence of a normal response at 40 dB was considered a moderate hearing loss; absence of a response at 40 and 60 dB a severe hearing loss; and an absent response at all three levels a profound hearing loss.





Figure 5. Pure-tone audiometry measured in children aged 5 and above.

Table 6. WHO hearing impairment grades, definitions, and recommendations.<sup>217</sup>

Grade of hearing impairment <sup>a</sup>	Clinical hearing	Recommendations
Slight (26–40 dB HL)	Normal voice at a distance of 1 metre	Counselling; hearing aids recommended for children with thresholds > 30dB
Moderate (41–60 dB HL)	Raised voice at a distance of 1 metre	Hearing aids recommended
Severe (61–80 dB HL)	Some words when shouted into better ear	Hearing aids; if not available, signing and lip-reading necessary
Profound or deaf (≥81 dB HL)	Unable to understand even with shouting into ear	Hearing aids may prove useful; lip-reading, signing, additional rehabilitation essential

<sup>a</sup> Defined as the average pure-tone threshold of 0.5, 1, 2 and 4 kHz in the better ear.

The BAEP screening was performed without sedation and under the supervision of at least one of the researchers. Older children were seated on a chair and those younger in their guardian's lap. Attempts were made to keep each child's movements, especially eye movements, to the minimum, and the youngest children



Figure 6. Brainstem auditory-evoked potentials registered without sedation.

were calmed by breastfeeding. After the skin was first carefully dried from sweat, the electrodes were positioned on the anterior forehead and at the mastoids. In the case of excessive artefacts in the response curve, the measurement was temporarily paused until the artefact source was neutralized.

## 7.5 Statistical analysis

Statistical analyses of the data were performed with Statview® (v.5.0.1, SAS Institute Inc., Cary, NC, USA). Statistical correlations for nominal variables were analyzed with Pearson's chi-square and Fisher's exact tests, whichever was feasible. Medians of continuous variables were compared with Mann-Whitney's U-test. Resulting p values  $\leq 0.05$  were considered statistically significant.

## 8. RESULTS

### 8.1 Respiratory infections – virology and clinical profiles [I]

In this study, of 102 NPS specimens, 37 (36%) appeared positive in PCR: 27 (26%) for HRV alone, 3 (3%) for HEV alone, and 7 (7%) for both viruses. In comparison of the virus-positive and -negative children, the virus-positive children were significantly younger (36 months vs. 52 months;  $p=0.024$ ). Chronic illnesses (HIV, SCD, tuberculosis) were present in 70% (26/37) of the PCR-positive vs. 63% (41/65) of the PCR-negative children ( $p=0.604$ ), and malnourishment in 19% vs. 31% ( $p=0.218$ ). No statistically significant differences emerged in frequencies of clinical findings among the PCR-positive vs. PCR-negative children with common colds: feverishness [12 (30%) vs. 13 (20%);  $p=0.161$ ], rhinorrhea [25 (68%) vs. 35 (54%);  $p=0.176$ ], cough [22 (59%) vs. 28 (43%);  $p=0.112$ ], AOM [9 (24%) vs. 8 (12%);  $p=0.117$ ], pneumonia [6 (16%) vs. 6 (9%);  $p=0.292$ ]. Neither did other clinical findings, such as wheezing, postnasal drip, tonsillitis, hoarseness, or conjunctivitis, associate with the PCR outcome, and these findings occurred in only a few children. Five of the six picornaviral cold-associated pneumonias occurred in HIV-positive children.

Both the occurrence throughout the study of common colds and the proportion of HRV-positive specimens increased, HRV presenting in 47% (25/53) of the specimens taken in the coolest month, July, vs. 18% (9/49) in April to June combined ( $p=0.021$ ; Figure 7).




Figure 7. Emergence of human rhinovirus (HRV) and human enterovirus (HEV) in polymerase chain reaction in nasopharyngeal smears of paediatric outpatients with respiratory symptoms during 4 autumn-winter months in Luanda, Angola. The monthly mean temperatures (left vertical axis) and air humidity values (right vertical axis) are long-term averages in Luanda. (216) Adapted from: Taipale A, Pelkonen T, Roivainen M, Kaijalainen S, Bernardino L, Peltola H, Pitkäranta A. Human rhino- and enteroviruses in children with respiratory symptoms in Luanda, Angola. Accepted for publication in Paediatrics and International Child Health; with kind permission from Maney Publishing ([www.maneypublishing.com/index.php/journals/pch](http://www.maneypublishing.com/index.php/journals/pch)).

## 8.2 Chronic suppurative otitis media – clinical and microbiological variables [II]

Background characteristics, medical history, and clinical findings were compared between 23 children with and 23 controls without CSOM (Table 7). CSOM was bilateral in 10 of the 23 (43%) cases, and its duration ranged from 2 months to 7 years (median 12 months). Six (26%) CSOM children suffered from both HIV and tuberculosis. None of the 23 children with CSOM exhibited intra- or extracranial complications.

The bacterial aetiology of CSOM appeared monomicrobial in all 18 bacterial-positive cases and comprised *Proteus* (8, 44%), *Pseudomonas* (4, 22%), *Shigella* (1, 6%), and *Haemophilus* (1, 6%) species. Four (22%) smears remained culture negative. Among these bacteria, antibiotic resistance was common against amoxicillin (64%), amoxicillin-clavulanic acid (60%), and trimethoprim-sulpha (71%), but no resistance occurred against ciprofloxacin or third-generation cephalosporins.

Among the CSOM children, slight-to-severe hearing loss (PTA or BAEP 28 to 80 dB HL) occurred in 52% (17/33) of the discharging ears, and slight hearing loss (PTA 28 to 31 dB HL) occurred in 23% (3/13) of the non-discharging ears, of which 2 had dry TM perforations and one a retracted TM. Bilateral hearing loss was present in 30% (7/23) of the CSOM children, and 6 children exhibited a PTA threshold > 30 dB HL in the better ear. Only one 8-year-old child had had attention paid previously to his hearing loss. None of the control children had bilateral hearing loss.

## 8.3 Sickle-cell disease-associated otorhinolaryngological pathology [III]

This study comprised 61 children with and 61 controls without SCD. Both groups included 36 (59%) boys, and the median ages of the children were 4.9 years (range 0.7-15.7) in the SCD and 4.8 (range 0.8-13.7) in the control group. Among the SCD children, 41 (67%) took regular medication (mainly folic acid, paracetamol, and ibuprofen), and 12 (19%) had received blood transfusions. For no SCD children were penicillin prophylaxis and pneumococcal vaccines available.

Table 8 shows that no significant differences occurred among the SCD vs. control children in ORL findings, including middle-ear and TM pathology.

In pure-tone audiometry screening, failure to hear a 25 dB HL stimulus at one or more frequencies was the case for 15 (60%) SCD- and 14 (50%;  $p=0.465$ ) control children. Bilateral hearing loss occurred in 9 (36%) SCD and 3 (11%;  $p=0.047$ ) control children (Figure 8). These hearing losses were of slight degree (30-40 dB HL), except for solitary losses of moderate degree in two SCD and two control children. None appeared to need a hearing aid. In SCD vs. control children, neither the average hearing thresholds nor the failure rates differed significantly at any frequency.

Table 7. Selected background features and clinical findings of children with (N=23) and without (N=23) chronic suppurative otitis media (CSOM) in Luanda, Angola. Comparisons by Pearson's chi-square and Fisher's exact tests. Adapted from: Taipale A, Pelkonen T, Taipale M, Bernardino L, Peltola H, Pitkäranta A. Chronic suppurative otitis media in children of Luanda, Angola. 2011; 100: 84-88. © 2011 The Authors / *Acta Paediatrica* © 2011 Foundation Acta Paediatrica.

	CSOM children n (%)	Controls n (%)	p value
<i>Background features</i>			
Female	11 (48)	11 (48)	
Median age (years); range	4.3; 1.1-11.5	4.5; 0.8-12.5	
Number of children in the household; range	2; 1-7	3; 1-7	0.583
Running water at home	8 (35)	6 (26)	0.749
Electricity <sup>a</sup> at home	16 (70)	16 (70)	>0.999
Attendance at national vaccination program	20 (87)	17 (74)	0.265
Median duration of breastfeeding, months (range)	18 (1-24)	18 (3-37)	0.114
Earlier episodes of otorrhoea	23 (100)	3 (14)	< 0.001
Ear pain, earlier or currently	23 (100)	0	< 0.001
<i>Clinical findings</i>			
Acute pulmonary infection	9 (39)	1 (4)	0.010
Cervical lymphadenopathy > 1cm	15 (65)	5 (22)	0.007
Tuberculosis	8/22 (36)	0	0.002
HIV positivity	14/22 (64)	0	< 0.001
Malnutrition	8 (35)	2 (9)	0.071

<sup>a</sup> During several hours 1-2 times daily through public electric transmission




Figure 8. Mean hearing thresholds in decibels (lines; left vertical axis), and the numbers of children failing to hear a 25 dB stimulus with their better ears (columns; right vertical axis), among 25 children with sickle-cell disease (SCD) and 28 controls in Luanda, Angola. Adapted from: Taipale A, Pelkonen T, Bernardino L, Peltola H, Pitkäranta A. Hearing loss in Angolan children with sickle-cell disease. *Pediatrics International* 2012; 54: 854-857. © The Authors. *Pediatrics International* © Japan Pediatric Society.

## Results

Table 8. Otorhinolaryngological profiles in 61 children with sickle-cell disease (SCD), their 61 controls, and 78 children with human immunodeficiency virus (HIV) and their 78 controls in Luanda, Angola (compared by Pearson's chi-square and Fisher's exact tests).

	SCD			HIV		
	Patients n (%)	Controls n (%)	p value	Patients n (%)	Controls n (%)	p value
<i>History</i>						
Otalgia	25 (41)	31 (51)	0.276	44 (56)	27 (35)	0.006
Otorrhea episodes	6 (10)	11 (18)	0.191	34 (44)	17 (22)	0.004
Hearing complaints	5 (8)	9 (15)	0.395	6 (8)	9 (12)	0.415
Speech delay	1 (2)	2 (3)	>0.999	2 (3)	1 (1)	>0.999
<i>Current findings</i>						
Respiratory infection	29 (48)	22 (36)	0.286	55 (71)	25 (32)	<0.001
Dermal lesions <sup>a</sup>	3 (5)	3 (5)	>0.999	25 (32)	4 (5)	<0.001
Oral candidiasis	1 (2)	3 (5)	0.619	9 (12)	5 (6)	0.402
Dental caries	26 (43)	23 (38)	0.580	44 (56)	25 (32)	<0.001
Tonsillar hypertrophy <sup>b</sup>	10 (16)	6 (10)	0.334	1 (1)	6 (8)	0.063
Salivary gland hypertrophy	2 (3)	0	0.496	2 (3)	0	0.497
Cervical lymph nodes <sup>c</sup>	11 (18)	4 (6)	0.096	35 (45)	8 (10)	<0.001
Acute otitis media	2 (3)	4 (6)	0.680	8 (10)	9 (12)	>0.999
Chronic suppurative otitis media	0	2 (3)	0.496	21 (27) <sup>d</sup>	3 (4)	<0.001
Middle-ear effusion	1 (2)	1 (2)	>0.999	0	0	>0.999
Dry tympanic membrane perforation	1 (2)	53 (48)	>0.999	7 (9)	1 (1)	0.063
Cerumen impaction	27 (44)	30 (49)	0.586	26 (33)	34 (44)	0.249

<sup>a</sup> Maculopapular eczema, folliculitis, molluscum contagiosum

<sup>b</sup> At least 50% of the space between pillars

<sup>c</sup> At least 1 cm

<sup>d</sup> Eight (38%) cases bilateral

Of the 15 SCD children (6 to 16 years old) who exhibited uni- or bilateral hearing loss, 2 children (3 ears) had cerumen impaction; of the rest, none manifested AOM, MEE, or TM perforations. Two children had previously noticed their own hearing loss.

#### **8.4 Human immunodeficiency virus-associated otorhinolaryngological pathology [IV]**

This study compared ORL status in 78 HIV-positive children and 78 controls. The median age in the HIV vs. control group was 4.3 years (range 0.8-14.8) vs. 4.1 years (range 0.8-13.7), and both groups included 36 (46%) boys. Of the HIV-positive children, 41 (53%) showed clinical signs of advanced or severe disease according to WHO criteria; 36 (46%) received antiretroviral treatment, 61 (78%) antibiotics (mainly sulfa-methoxazole), and 5 (6%) anti-tuberculosis treatment. Concurrent or previous diagnosis of tuberculosis occurred in 22/77 (29%) HIV-positive children vs. 2/77 (3%;  $p<0.001$ ) controls, current diagnosis of pneumonia in 17 (22%) HIV-positive vs. 2 (3%;  $p<0.001$ ) controls, and current hepatomegaly in 8 (10%) HIV-positive vs. zero controls ( $p=0.007$ ).

Major differences emerged in the ORL findings among the HIV-positive vs. control children, as described in Table 8. None of the clinical variables inside the HIV group were associated with these children's antiretroviral treatment status.

Hearing screening by BAEP was performed in all study children, and 29 HIV-positive and 31 control children also underwent audiometry. No data on hearing could be obtained in 2 young control children (3 ears), in whom BAEP screening failed, and pure-tone audiometry was nonfeasible. After combining the results from these two screening methods, 20 (26%) HIV-positive and 12 (15%) control children exhibited slight-to-profound hearing loss. Bilateral hearing loss was present in 10 (13%) HIV-positive children vs. one (1%;  $p=0.009$ ) control child. In the HIV-positive children, otoscopy of the 30 hearing-impaired ears revealed CSOM (12; 40%), cerumen impaction (7; 23%), dry TM perforation (5; 17%), intact, mobile TM (4; 13%), and retracted TM (2; 7%). Hearing defects among the HIV-negative children (13 ears) related to cerumen (5; 38%), to unknown factors (5; 38%), to AOM (2; 15%), and to CSOM (1; 8%).



## 9. DISCUSSION

### 9.1 The role of picornaviruses in tropical respiratory infections

In the present study, screening picornaviruses in children with respiratory symptoms from various outpatient clinics, the incidence of respiratory symptoms increased in the cool season. Picornaviruses occurred at young ages, a median 3 years, with symptoms of fever, rhinorrhea, and cough. The frequencies of these symptoms, as well as otitis media (discussed later in detail) and pneumonia, showed no alteration related to the children's picornavirus positivity or underlying chronic conditions. Furthermore, neither HIV infection, SCD, nor malnutrition associated with picornavirus positivity.

In this sample, proportions of children with HIV infection and SCD were highly overrepresented in relation to those of the general population. This is largely due to the high contribution of these chronic illnesses to health care: In the mid-2000's, systematic HIV testing of patients in the Paediatric Hospital of Luanda revealed roughly one-fifth to be HIV positive.<sup>216</sup> Notably, the present study's SCD and HIV children exhibited no higher frequencies of acute respiratory symptoms than in control children.

Here, prevalence rates for HRV and HEV positivity were consistent with those in other recent studies conducted among outpatients in subtropical or tropical countries.<sup>48,49,51,217,218</sup> Strictly speaking, the present study could verify no causality between viral shedding and clinical infection, as that would have demanded serial NPS sampling, and obtaining and processing multiple samples from the same individuals would have posed excessive economic and logistical challenges regarding the study context.

Our findings suggest that HRV transmission may increase in the tropics at decreasing temperatures: HRV caused nearly half of all RTI cases in the coolest month of July. However, the limited study period (April to July), as well as the relatively small sample size, restricts further speculation on picornavirus seasonality. None of the larger-scale studies from tropical regions reported a clear seasonality for picornaviruses: association with weather parameters went either undetectable<sup>48,49</sup> or unanalyzed.<sup>217</sup> A cohort study from tropical Brazil perceived a peak in HRV incidence during the local autumn, however.<sup>50</sup>

Knowledge of the aetiology and epidemiology of RTIs is recognized to be of major importance in developing countries where these diseases carry high morbidity and mortality.<sup>219</sup> Until recently, HRV has been considered a rather harmless virus responsible mainly for mild RTI symptoms, and it has gained little attention in research in developing countries. Because up-to-date evidence links HRV not only



to lower RTIs,<sup>21,22,33,48,53,220-222</sup> but also to invasive pneumococcal diseases,<sup>223,224</sup> and even to meningitis,<sup>39</sup> research on picornavirus infections in developing countries becomes more relevant. In addition, the underlying nutritional or immune deficiencies and their impact on the clinical picture and outcomes of common cold deserve more attention.

Because the multiplicity of HRV and HEV types hampers effective vaccine development, restricting the transmission of these viruses remains the major preventive measure.<sup>219</sup> Further research is therefore needed on the epidemiological and clinical aspects of HRV and HEV infections in developing countries.

## **9.2 Middle-ear pathology**

### **9.2.1 OTITIS MEDIA AS A COMPLICATION OF RESPIRATORY INFECTION**

AOM occurred in one-quarter of the children with picornavirus infections. More than half of all AOM children who underwent NPS sampling actually yielded picornavirus, but this association lacked statistical significance. Should any real association between AOM and picornaviruses among all RTI-causative agents have existed, demonstrating it would probably have demanded a considerably larger sample size, because of the high overall frequency of AOM during viral RTIs. As such a high proportion of children with RTI will develop AOM anyway, picornaviruses may not, from an epidemiological point of view, outweigh other aetiopathological viral agents of AOM in their middle-ear invasiveness but simply by their higher frequency. Actually, the proportions of AOM and picornaviral RTI that we obtained agree with findings from temperate climates, where picornavirus RTIs lead to AOM in approximately one out of three cases,<sup>18</sup> and were present in up to half the AOM children.<sup>41,42,57,58</sup>

This study suggests a marked involvement of picornaviruses in the AOM complication of tropical common colds. The extent of this involvement remains a subject for further research in the quest for more accurate characterization of AOM – the major pathway to CSOM in areas of a high burden of chronic otitis media.

### **9.2.2 CHRONIC SUPPURATIVE OTITIS MEDIA**

We evaluated CSOM in children with otorrhoea of up to 7 years' duration. Studies from developing countries verify that children may suffer from extremely longstanding otorrhoea, because treatment is delayed or lacking.<sup>70,81,111,112,225</sup> The median age of our CSOM children was 4 years. Several earlier studies corroborate the young age of CSOM patients.<sup>65,73,112,115,116,225</sup> As for ours, more than 40% had bilateral disease, similar to common findings in childhood CSOM in developing countries.<sup>70,73,80,82,112,225</sup> All of these features – long duration, early onset, and a tendency toward bilaterality

– underline the importance of prompt treatment of CSOM during the child’s critical age of linguistic and cognitive development.

The present study excluded children whose otorrhoea had persisted for less than 2 months. Worldwide, the definition of CSOM is not uniform, and durations of at least 2 weeks to 3 months have been suggested as defining middle-ear discharge as being chronic. In the Paediatric Hospital of Luanda, the vast majority of children with otorrhea had at least several months’ history of the disease, and thus the data obtained by applying the cut-off limit of 2 months represented the actual patient material well.

Our most frequent bacterial causative agents were *Proteus* and *Pseudomonas* species. This is indeed typical of developing countries,<sup>65,66,69-71,73,112</sup> except that *Staphylococcus aureus* was absent from our patients. However, our sample size was limited, and therefore coincidence most probably explains this non-emergence of *Staphylococci*. Additionally, pus sampling from the ear canal, instead of as an aspirate directly from the middle ear, may have led to a higher risk of contamination. On the other hand, one study from India detected no difference between these two sampling methods, and so our results may be considered rather reliable.<sup>226</sup>

The high frequency of opportunistic Gram-negative rods reflects the poor general state of health and the prolonged disease process in our CSOM children and thus points to their condition’s severity.

Mycobacterium diagnostics, had it been performed, would have provided valuable additional information because pulmonary tuberculosis occurred in more than one-third of our CSOM children. Considering this high tuberculosis prevalence, as well as the severity of hearing loss in some patients, some of our children may actually have suffered from aural tuberculosis. In general, tuberculous otorrhea is regarded as a rare disease offering diagnostic challenges.<sup>227,228</sup> In future, a study on mycobacterial CSOM among HIV-positive children in resource-poor settings would be of value.

No intra- or extracranial complications of CSOM were encountered. This was rather expected, as the CSOM children were outpatients visiting the clinic for other reasons. However, because complications of CSOM continue to cause morbidity and mortality in children of sub-Saharan Africa, the subject requires research in the settings where such cases more likely will emerge, i.e. in ORL clinics or during acute health care. For the same reason – CSOM-induced adverse outcomes – general research on childhood CSOM in developing countries needs to continue.

### 9.2.3 MIDDLE-EAR PATHOLOGY IN SICKLE-CELL DISEASE

Regarding the impact of SCD on childhood middle-ear infections in Luanda, unexpectedly, SCD children only occasionally exhibited middle-ear pathology, despite their lack of penicillin prophylaxis and pneumococcal vaccines. Few reports

concerning otitis media in SCD children exist: 3 hearing-screening studies (2 from the USA, one from Nigeria), reported MEE in up to 27.5%.<sup>139,142,159</sup> Of these, one US study perceived an increased prevalence of middle-ear pathology in SCD vs. control children,<sup>142</sup> while in Nigeria the rates appeared equal.<sup>139</sup> In addition, a retrospective survey from the USA detected fewer tympanostomy-tube insertions in SCD children than in a historical SCD-free cohort.<sup>229</sup> Recurrent, persistent, and complicated episodes of otitis media are, however, characteristic of children without haemoglobinopathies but with antibody deficiencies.<sup>165,230,231</sup> SCD children with their weakened splenic function might therefore be expected to show a tendency towards acute and chronic otitis and its severe suppurative complications, but actual evidence on the subject remains small and controversial.

Possible explanations for the absence of middle-ear pathology among our SCD children include a better clinical balance in this group, when compared to the worse-off majority not followed up at the SCD clinic; survival of children with a milder form of SCD (the most severe cases leading to early mortality); and, finally, mere coincidence. Concerning this study, we must also emphasize the possibility of missing small amounts of MEE in young children with suboptimal otoscopic views due to impacted cerumen in the ear canal. Future studies should include a larger and wider sample of SCD children and controls, as well as follow-up.

Some studies suggest only a low proportion of *Streptococcus pneumoniae* in bacteraemia of SCD children in malaria-endemic regions; this has raised the question whether malaria-induced splenomegaly could partially preserve splenic function in these children.<sup>232,233</sup> Malaria may thus be one of the environmental factors modifying the course of SCD. A follow-up study is required to verify the impact of SCD on the incidence and course of childhood otitis media in low-resource, malaria-endemic regions where both SCD and otitis-related deaths occur.

#### 9.2.4 THE BURDEN OF OTITIS MEDIA IN HUMAN IMMUNODEFICIENCY VIRUS-POSITIVE CHILDREN

In all HIV-positive children of the present study, CSOM was present in more than one-quarter and dry perforations in one-tenth. Our small and hospital-based samples may have been biased, and our CSOM group probably exaggerates the true contribution of HIV in CSOM epidemiology. However, several studies confirm the high burden of chronic otitis in HIV-positive children,<sup>201-204</sup> and reports also confirm the occurrence of systemic complications of otitis media among them.<sup>187,201</sup>

No previous reports were available of incidence rates or wider statistics concerning the role of HIV in otitis media and its complications in developing countries. In Brazilian HIV-positive children, management of HIV alleviated the burden of their otitis media.<sup>207</sup> The impact of the HIV epidemic on the course and consequen-

ces of CSOM is probably considerable in regions of high HIV prevalence, and thus deserves future research.

### 9.3 Otorhinolaryngological profiles in different patient groups

Of every ten of our children without clinical suspicion of HIV, seven exhibited ORL pathology, making ORL pathology apparently highly frequent. The most typical ORL symptoms and diseases were respiratory infections, lymph-node enlargement, ear pain, acute or chronic otitis media, and oral thrush. In children of developed countries, RTIs<sup>234</sup> and AOM<sup>235</sup> are also usual. However, chronic otitis and oral candidiasis are ORL presentations typical of children who experience poor living conditions and nutrition and hygiene deficiencies.

In our children, cerumen impaction was frequent. The children or their caregivers were only rarely familiar with the meaning or normal function of ear wax. Indeed, cerumen impaction is common in children of resource-poor countries and may even lead to clinically significant hearing loss.<sup>236-239</sup> In children with already compromised hearing due to other causes, even slight additional cerumen-induced hearing loss may significantly worsen hearing ability. In general, not many studies concern the adverse effect of excessive cerumen on hearing. Among some adults in the UK, one-third benefited from removal of impacted cerumen, and even though the procedure improved hearing levels by an average of only 5 dB HL, some individuals' hearing levels improved by up to 36 dB HL.<sup>240</sup>

In populations with high prevalences of TM perforations, limited knowledge of anatomy of the ear, and poor availability of clean water or cerumenolytic agents, removal of cerumen should occur in health care centres with trained personnel, manual removal without irrigation probably being the most feasible method.

In this study, of every ten HIV-positive children, no less than nine exhibited ORL pathology. This is consistent with reports that also uniformly suggest substantial ORL involvement in paediatric HIV infection.<sup>184-186,188</sup> Compared to our controls, the present study's HIV-positive children presented a higher number of pathologic findings with a more severe clinical picture. They had marked cervical lymphadenopathy and rampant caries more frequently, but so did some control children, as well. In the head and neck, bilateral CSOM and widespread eczema appeared particularly strong indicators of HIV positivity. Compared to earlier reports, we detected HIV-associated oral candidiasis somewhat less and parotid enlargement much less frequently.<sup>184-186,188</sup> On the whole, our findings suggest that, as part of their medical follow-up, HIV-positive children should routinely undergo an ORL examination.

In contrast to the HIV patients, the present study was unable to demonstrate any significant elevations in the frequencies of ORL findings in the SCD children. No other comparative studies on the subject have emerged. The present study's SCD

children also exhibited none of the reported ORL complications of SCD, such as deep infections, osteomyelitis, or marked tonsillar hypertrophy.<sup>166,170</sup> This suggests that such findings are fairly unusual even in developing countries. However, because ORL complications of SCD do exist in case reports, defining their occurrence would be interesting in resource-poor settings where the majority of the patients live.

## **9.4 Hearing Loss**

### **9.4.1 HEARING LOSS DUE TO CHRONIC SUPPURATIVE OTITIS MEDIA**

The present study detected at least slight hearing loss in half the CSOM ears and bilateral hearing loss in nearly one-third of CSOM children. Only one of these cases had been noticed by the family. These characteristics are consistent with findings in developing countries that of every 10 CSOM children, hearing loss occurs in 5 to 9.<sup>79-82,114-116</sup> Data also show that both otorrhoea and hearing loss in children in resource-poor settings remain widely neglected.<sup>73,112,241</sup> Unfortunately, the onset of CSOM during infancy may pose a higher risk for hearing loss.<sup>116</sup> Chronic otitis media occurring at school age may cause hearing loss sufficiently severe to endanger scholastic performance.<sup>82,117,239</sup> Therefore, in resource-poor communities, more precise recognition of a child's ear discharge as a warning sign is warranted in order to fully utilize existing resources for prevention of and rehabilitation of hearing loss.

Information should be targeted to communal workers serving children and families at clinics and schools, as well as to primary health care professionals. As CSOM children typically suffer from slight-to-moderate hearing loss, they would, even without access to hearing aids, be substantially supported by simple rehabilitative measures such as having seats in the front of the classroom, being addressed in a loud and clear voice, and making lip-reading more possible – instead of their being erroneously regarded as lazy or unintelligent.

### **9.4.2 HEARING LOSS IN SICKLE-CELL DISEASE**

In audiometric evaluation of our SCD children, screening failures occurred in three-fifths of them and bilateral hearing loss in more than one-third. However, only bilateral hearing loss was significantly associated with SCD. The absence of evidence of middle-ear pathology suggests that the hearing impairment may have originated mainly from the inner ear. Of the 15 hearing-impaired SCD children, however, in 2 (3 ears), cerumen impaction prevented reliable otoscopic and tympanometric evaluation and may have been the actual reason for compromised hearing. Only 2 hearing-loss cases had previously been detectable. None presented with a PTA threshold >30 dB HL in the better ear, the current WHO standard for considering a hearing aid.<sup>4</sup> The unexpectedly high hearing-loss rate among the present study's

control children may have been coincidental, or a result of higher participation of families of children with ear- or hearing problems.

We detected a relatively high rate of hearing loss compared to that for previous researchers, particularly if losses attributable to middle-ear pathology in the earlier data are excluded. Notably, half the hearing-impaired children were only 5 to 8 years of age, younger than in most studies, which predominantly examined older children.<sup>139,142-145</sup> We thus highlight the risk for hearing loss in SCD that occurs even as early as at ages 5 to 6. In contrast to some reports,<sup>142,145</sup> we found no severe impairment, which suggests that severe hearing loss occurs in children only rarely. However, even slight or fluctuating hearing loss may adversely affect a child's speech development.<sup>242</sup>

With improved management, more SCD children will survive in sub-Saharan Africa beyond early childhood.<sup>119</sup> In these survivors, screening for hearing problems should be implemented in their routine care even at preschool age.

#### **9.4.3 HEARING PROBLEMS IN HUMAN IMMUNODEFICIENCY VIRUS-POSITIVE CHILDREN**

Among the HIV-positive children, some degree of hearing loss occurred in one-quarter, and half of those suffered bilateral loss. More than half the hearing loss occurred in ears with TM perforations, the majority of which were actively discharging. This is consistent with data in other studies from developing countries, showing hearing loss in 23 to 39%.<sup>191-194</sup> These studies also frequently detected middle-ear pathology behind the impairment. In resource-rich settings, the proportion of otitis media in HIV-associated hearing loss may be lower, but risk for hearing loss remains elevated.<sup>243</sup>

The present study confirms and underlines the importance of recognizing hearing loss and treating otitis media promptly in HIV-positive children in developing countries, as well as the importance of screening their hearing on a regular basis.

### **9.5 Practical and ethical viewpoints**

Our study was conducted in a paediatric hospital in tropical Angola, where few data exist on childhood ORL problems. The study children were simply recruited among the hospital's outpatients, which provides both strengths and limitations for the study. Obviously, the samples were highly biased compared to the general population: In addition to chronic underlying diseases, higher socioeconomic classes may have been overrepresented among the hospital's outpatient population, as even public transportation to the hospital may have been too expensive for the poorest families. Moreover, factors such as higher education, positive family history, or

suspicion of existing ORL problems may have furthered the guardian's decision to participate in the study. However, in developing countries, children ask for medical care due to ORL problems only rarely. These problems are and should therefore be screened for and detected during visits for other reasons; this underlines the clinical relevance of the present findings.

In resource-poor Angola, basic ORL services, even ones such as otoscopic examination, are available only rarely. Soundproof facilities for hearing screening and ear-suction devices for cleansing the ear canal were nonexistent, and electricity was undependable. These circumstances contribute to this study's limitations: pure-tone thresholds obtained, especially in the lowest frequencies, may be somewhat higher than in reality, and in some cases (especially in young children) not all cerumen could be removed from the ear canal. BAEP screening proved challenging due to excessive sweating when the air-conditioner was broken or lacked electricity. Such circumstances represent, however, the usual settings in health care and research practiced in resource-poor countries.

Pure-tone thresholds were measured without masking the contralateral ear. Due to transaural bone conductivity, this may have resulted in erroneously low pure-tone thresholds in the worse-hearing ears in children whose pure-tone thresholds differed by more than 50 dB HL. In addition, as no bone-conduction thresholds were measured, a distinction between conductive and sensorineural hearing loss was impossible. Furthermore, because in cochlear hearing loss the correlation between hearing thresholds obtained by click-evoked BAEP vs. pure-tone audiometry is strong only at 2 to 4 kHz and remains weaker at lower frequencies,<sup>244</sup> the ability of our BAEP measurements to detect such hearing loss cases was probably suboptimal.

To be ethical, research in a resource-poor environment should have a clear purpose,<sup>245,246</sup> which, in the present study, was to collect epidemiological and clinical data on childhood ORL diseases affecting millions of children worldwide. In addition, making medically purposeful research ethically sustainable requires ethical evaluation by local professionals, as well as consent from the study attendants. The present study underwent ethical consideration in the Paediatric Hospital of Luan-da.<sup>245,246</sup> Because of limited literacy among Angolans, however, study information and consent to participate was handled orally. The majority of the guardians were familiar with neither the usual childhood ORL diseases nor the concept of medical research.

According to Ruth Macklin, these examples represent a true dilemma for research among resource-poor communities: should ethical guidelines arise from pragmatism or idealism; how can one create guidelines that are, as Macklin noted, "usefully prescriptive without being hopelessly aspirational"? In a culture which knows no basic concepts of medicine or medical research, how to adapt research procedures, such as obtaining informed consent, to the environment in such a way that the consent obtained becomes genuine? Macklin states that research concerning a major health issue in the target population can be ethically justified, if no



possibility exists for conducting it in another population with more potential for understanding its content.<sup>247</sup> In the team of our present study, the local professionals plus one member with long-term local experience lowered these cultural barriers.

In addition to the wider advantages for future patients, short-term research may locally provide for the participants “ancillary care” not directly related to the study protocol.<sup>245,246</sup> Here, ancillary care (e.g. medical prescriptions) was provided when needed, but did not serve as a “decoy” in recruitment, although sometimes families with known ear complaints appeared more willing to participate in hope of free treatment. Leaving such families virtually without access to any treatment would have been the most unethical option. Unfortunately, in the absence of any local ORL service network, no follow-up for these children was available.

In Luanda, resources for rehabilitation of the hearing impaired are poor. Some study participants could be advised to contact a local school for the hearing impaired, but hearing-aid rehabilitation was unavailable. However, in resource-poor communities, even recognizing the hearing impairment would help the child’s family and teachers to support the child in daily communication.<sup>236</sup>

ORL diseases and hearing loss in children of developing countries are largely neglected areas in medical research, with more data required. Worldwide, 90% of the need for hearing aids exists in developing countries, where only less than one in 40 will receive the required services.<sup>4</sup>

In resource-poor countries, vast, unselective screening programs requiring expensive equipment and training of personnel may be unfeasible in the absence of any functioning referral and rehabilitation system. However, not to screen at all would inevitably lead to permanent disability for hundreds of thousands of hearing-handicapped children.<sup>248</sup> More regional data on childhood hearing loss is therefore essential for targeted screening.<sup>7,14,249</sup> Questionnaires are one tool proposed for identifying children in need of hearing screening. The present findings suggest that, should such questionnaires be applied, they should include the child’s HIV and possibly also SCD status. Additional simple methods for hearing assessment in primary health care, such as whispering tests,<sup>17,250</sup> also need research and validation in developing countries.

The child’s optimal cognitive and social development relies on adequate hearing, and underprivileged communities need the full potential of their children, the builders of the future. The lifetime cost of childhood hearing loss has been estimated to exceed that of the cases with later onset.<sup>7</sup> In resource-poor settings, therefore, prevention and rehabilitation of hearing loss generally deserve higher priority, and the highest priority for these services should belong to children.



## 10. SUMMARY AND CONCLUSIONS

A massive shortage of ORL services, education, and research exists in developing countries where the resource-poor environment poses risk factors for chronic and recurrent ORL diseases. The burden of ORL diseases remains therefore high in these areas among children. In sub-Saharan Africa, one ORL expert may serve millions of people, and awareness of ORL diseases in primary care remains limited. Hearing loss and its impact on society are under-recognized, and only one in 40 in need of hearing rehabilitation receives the needed device. In the children of Africa, the HIV epidemic and high prevalence of sickle-cell disease and tuberculosis further affect the pathology and conditions related to the ORL area.

The present study was conducted in the Paediatric Hospital of Luanda, Angola, to collect data on ORL pathology and hearing in children of various outpatient clinics, and particularly to define the occurrence of HRV and HEV during the cool season, the clinical and microbiological characteristics of childhood CSOM, and the HIV- and SCD-associated ORL findings and hearing loss. Background features and medical history came from the children's guardians, and the children underwent general physical and ORL examination, plus hearing screening by BAEP and pure-tone audiometry.

Picornaviruses were evident in more than one-third of common-cold patients. Common-cold symptoms, as well as the proportion of rhinovirus as the causative agent, increased as the temperature fell. The virus-positive children were significantly younger than the virus-negative, and AOM complicated the picornaviral RTIs in about one-quarter. In resource-poor living conditions, chronic ear discharge typically harbored gram-negative rods, and in the setting of outpatient clinics, the vast majority of the patients suffered from HIV or tuberculosis; approximately one-third had slight-to-severe bilateral hearing loss. SCD children also presented with an ascending risk for hearing loss, mainly in the low and middle frequencies; these losses appeared slight, and the children exhibited no signs of the alleged susceptibility to otitis media. Abnormal ORL findings such as cervical lymphadenopathy, RTI, and otitis media were common in all children, but the children with HIV infection appeared to be major carriers of the burden of ORL diseases. Moreover, hearing loss was associated with HIV and mainly coincided with middle-ear pathology; the present study did not, however, rule out concomitant inner-ear pathology.

Several conclusions can be drawn. This work suggests the marked involvement in sub-Saharan Africa of picornaviruses, particularly HRV, in AOM epidemiology. In order to prevent more severe forms and bacterial complications of viral RTIs, more complete understanding is needed for the epidemiology and clinical consequences of picornaviruses in tropical areas with low resources and high co-morbidity.

In developing countries, a child may grow up with chronic ear discharge and hearing loss without the guardians or even medical professionals recognizing the condition. Especially in areas of the HIV epidemic, the prevalence and epidemiology of CSOM require updating.

SCD may lead to hearing loss, and children aged 5 to 6 should already be undergoing audiometric screening. However, the incidence and clinical picture of otitis media remains unclear among children with SCD in sub-Saharan Africa, and deserves a prospective follow-up study.

Ear discharge, extensive rashes, and persistent cervical lymphadenopathy may be signs of underlying HIV infections, and follow-up of HIV-infected children should include thorough ORL status, hearing screening, and prompt treatment of middle-ear infections.

Focusing attention on children's ears and hearing could substantially benefit many underprivileged communities. Both from the perspective of humanity and economy, these children are too valuable a resource to be wasted through preventable disability and suffering.

## 11. ACKNOWLEDGMENTS

This work was carried out in the years 2008 - 2013 at the Institute of Clinical Medicine, Faculty of Medicine, University of Helsinki; at the Department of Otorhinolaryngology at Helsinki University Central Hospital; at the National Institute for Health and Welfare; and at the Hospital Pediátrico David Bernardino in Luanda, Angola.

I wish to express my gratitude towards the leader of this study and my supervisor, Professor Anne Pitkäranta, a person with great professionalism, optimistic determination, and clear-minded wisdom. I felt privileged to work under her practical, straightforward, and encouraging guidance.

I would also like to express my appreciation to Docent Heikki Rihkanen, the Head of the Otorhinolaryngology Department, and Docent Antti Aarnisalo, Head of the Audiology Department, for the opportunity to work at the Department of Otorhinolaryngology. They also donated, from their departments, recycled devices on which the clinical execution of this study depended. In addition, I sincerely acknowledge Docent Merja Roivainen at the National Institute of Health and Welfare for her collaboration.

I owe respectful thanks to the official reviewers of this thesis: Docents Aino Ruohola from the University of Turku and Juha-Pekka Vasama from the University of Tampere. The accuracy of the reviewers resulted in marked improvements in my work, which Carol Norris PhD kindly revised for language despite her busy schedule.

My heartfelt gratitude goes to Tuula Pelkonen, MD PhD, for her hospitality during our stay in Angola in 2008. Her help was crucial in virtually everything. I also thank her for her friendship, co-authorship, and technical advice concerning writing and statistics. I sincerely thank Dr. Luis Bernardino, head of the Paediatric Hospital of Luanda and one of the designers of the study, for providing facilities and various forms of practical support. His efforts for the children of Luanda are invaluable. My thanks also go to our flexible assistant, paediatric nurse Dona Fátima, as well as to our gentle Angolan babysitter Madeleine Kotele.

I want to express my thanks to Professor Emeritus Heikki Peltola, whose meningitis trial in the Paediatric Hospital of Angola served as a platform for initiating our project. In addition, Irmeli Roine MD deserves acknowledgment for co-authorship, and laboratory researcher Svetlana Kaijalainen for analysis of virus samples. I would also like to thank my colleagues and co-workers in the audiophoniatic ward in the Department of Otorhinolaryngology, as well as my friend Veera Bergman, for their timely encouragement.

My deepest gratitude belongs to my family: my dear husband and companion Marko and our beloved children Siiri, Aaro, Hilma, and Vuokko. Marko also participated in data collection. I thank my parents Ulla and Juha Solasaari for their

lifelong support, and along with them my parents-in-law Salme and Pauli Taipale for their kind aid with babysitting.

Finally, I thank God for the hope which made this project purposeful. I travelled to Angola in a conscious search for hope, and, while visiting a church clinic in the rural south, happened to receive a gift from a local nurse: a t-shirt with a cross and two words in Kwanyama printed on it. Indeed, human life is worth cherishing: “*Otu Netelelo*” – “*We have hope*”.

This work received financial support from the Väinö and Laina Kivi Foundation, the Ear Research Foundation, the Pediatric Research Foundation, and research funds of the University of Helsinki.

Hyvinkää, August 2013

Anni Taipale

## KIITOKSET

Tutkimus toteutettiin vuosina 2008 – 2013 Helsingin yliopiston lääketieteellisen tiedekunnan kliinisellä laitoksella, Helsingin yliopistollisen keskussairaalan Korvaklinikalla, Terveiden ja hyvinvoinnin laitoksella sekä angolalaisessa Luandan lastensairaalassa (Hospital Pediátrico David Bernardino).

Haluan ilmaista kiittolisuuteni tutkimuksen johtajalle sekä väitöskirjatyöni ohjaajalle; erittäin ammatillisesti, optimistisen päämäärätietoisesti sekä selkeäpiirteisellä viisaudella toimineelle professori Anne Pitkärannalle. Hänen käytännönläheisessä, suoraviivaisessa ja rohkaisevassa ohjauksessaan työskenteleminen tuntui etuoikeudelta.

Haluan myös ilmaista arvostukseni Korvaklinikan johtajaa, dosentti Heikki Rihkasta, sekä audiologisen yksikön johtajaa, dosentti Antti Aarnisalaa kohtaan heidän antamastaan mahdollisuudesta työskennellä Korvaklinikalla. Lisäksi heidän osastoiltaan lahjoittamansa kierrätetyt tutkimuslaitteet mahdollistivat tutkimuksemme kliinisen toteutuksen. Kiitän myös lämpimästi yhteistyöstä dosentti Merja Roivaista Terveiden ja hyvinvoinnin laitokselta.

Ilmaisen kunnioittavat kiitokseni väitöskirjani esitarkastajille, Turun yliopiston dosentti Aino Ruoholalle sekä Tampereen yliopiston dosentti Juha-Pekka Vasamalle. Heidän tunnollisuutensa johti selvästi parempaan lopputulokseen työssäni, jonka kielentarkistuksen FT Carol Norris ystävällisesti otti tehtäväkseen kiireisestä aikataulustaan huolimatta.

Kiitän sydämellisesti LT Tuula Pelkosta hänen vieraanvaraisuudestaan viettäessämme aikaa Angolassa vuonna 2008. Hänen kaikkinaisen apunsa oli välttämätöntä. Olen hänelle kiittollinen myös ystävydestä sekä avusta artikkelien kirjoittamisessa ja tilastollisten menetelmien käytössä. Kiitän vilpittömästi Luandan lastensairaalan johtajaa tri Luis Bernardinoa, joka osallistui tutkimuksemme suunnitteluun, järjesti tilat tutkimuksen toteuttamiseen sekä tarjosi monenlaista käytännön apua. Hänen vaivannäkönsä Luandan lasten hyväksi on mittamattoman arvokasta. Haluan myös kiittää joustavaa tutkimusapulaistamme, lastensairaalan hoitajaa Dona Fátimaa sekä hellää angolalaista lastenhoitajaamme Madeleine Kotélea.

Haluan ilmaista kiittolisuuteni emeritusprofessori Heikki Peltolalle, jonka ohjaama aivokalvontulehduksen hoitoa koskeva tutkimus Luandan lastensairaalassa toimi alkuunpanijana omalle projektillemme. Lisäksi LT Irmeli Roine ansaitsee tunnustuksen osallistumisestaan artikkelin kirjoittamiseen ja laboratoriotutkija Svetlana Kaijalainen osallistumisestaan virusnäytteiden analysointiin. Haluan myös mainita kollegani ja työtoverini Korvaklinikan audiofoniatriiselta osastolta sekä ystäväni Veera Bergmanin, jotka rohkaisivat minua juuri oikeaan aikaan.

Syvin kiitollisuuteni kuuluu perheelleni: rakkaalle aviomiehelleni ja kumppanilleni Markolle sekä lapsillemme Siirille, Aarolle, Hilmaalle ja Vuokolle. Marko osallistui myös tutkimusdatan keräämiseen. Vanhempiani Ulla ja Juha Solasaarta haluan kiittää heidän elämänmittaisesta tuestaan, ja yhdessä heidän kanssaan kiitän appivanhempiani Salme ja Pauli Taipaletta avusta lastenhoidossa.

Lopuksi tahdon kiittää Jumalaa toivosta, joka antoi merkityksen työlleni. Matkustin Angolaan tietoisesti etsimään juuri toivoa, ja vieraillessani etelässä kirkon maalaisklinikalla satuin saamaan paikalliselta hoitajalta muiston: t-paidan, johon oli painettu ristin kuva sekä kaksi kwanyamankielistä sanaa. Ihmiselämä on todellakin vaalimisen arvoista; *”Otu Netelelo”* – *”Meillä on toivo”*.

Tämä työ sai rahallista tukea Väinö ja Laina Kiven Säätiöltä, Korvatautien tutkimussäätiöltä, Lastentautien tutkimussäätiöltä sekä Helsingin yliopiston tutkimusvaroista.

Hyvinkäällä elokuussa 2013

Anni Taipale

## 12. REFERENCES

1. Fagan JJ, Jacobs M. Survey of ENT services in Africa: need for a comprehensive intervention. *Glob Health Action* 2009; **2**(1): 1-7.
2. Fagan JJ. Developing World ENT: A global responsibility. *J Laryngol Otol* 2012; **126**(6): 544-547.
3. World Health Organization. Prevention of hearing impairment from chronic otitis media. Report of a WHO/CIBA foundation workshop, London: 19-21 November 1996. Geneva: World Health Organization; 1996.
4. World Health Organization: Guidelines for Hearing Aids and Services for Developing Countries. Geneva: World Health Organization; 2004.
5. CIA - The World Factbook 2012. The Central Intelligence Agency. (Accessed 10/10, 2012, at <https://www.cia.gov/library/publications/the-world-factbook/>).
6. McPherson B. Audiology: A developing country context. In: McPherson B, Brouillette R, eds. *Audiology in Developing Countries*. New York: Nova Science Publishers, Inc.; 2008: 5-20.
7. Olusanya BO, Newton VE. Global burden of childhood hearing impairment and disease control priorities for developing countries. *Lancet* 2007; **369**(9569): 1314-1317.
8. Najim O, Powell S. So you want to be an ENT surgeon? *BMJ Careers* 2007; 334: 205-207.
9. McPherson B, Olusanya BO. Screening for hearing in developing countries. In: McPherson B, Brouillette R, eds. *Audiology in Developing Countries*. New York: Nova Science Publishers, Inc., 2008:75-105.
10. World Health Organization. Chronic suppurative otitis media: Burden of illness and management options. Geneva: World Health Organization; 2004.
11. Brouillette R. Rehabilitation of hearing loss: Challenges and opportunities in developing countries. In: McPherson B, Brouillette R, eds. *Audiology in Developing Countries*. New York: Nova Science Publishers, Inc.; 2008:141-153.
12. Olusanya B. Early detection of hearing impairment in a developing country: what options? *Audiology* 2001; **40**(3): 141-147.
13. Olusanya BO. Early hearing detection at immunization clinics in developing countries. *Int J Pediatr Otorhinolaryngol* 2006; **70**(8): 1495-1498.
14. Gell FM, White EM, Newell K, Mackenzie I, Smith A, Thompson S. et al. Practical screening priorities for hearing impairment among children in developing countries. *Bull World Health Organ* 1992; **70**(5): 645-655.
15. Newton VE, Macharia I, Mugwe P, Ototo B, Kan SW. Evaluation of the use of a questionnaire to detect hearing loss in Kenyan pre-school children. *Int J Pediatr Otorhinolaryngol* 2001; **57**(3) :229-234.

16. Samelli AG, Rabelo CM, Vespasiano AP. Development and analysis of a low-cost screening tool to identify and classify hearing loss in children: a proposal for developing countries. *Clinics (Sao Paulo)* 2011; **66**(11): 1943-1948.
17. Clark JL, Newton V. Providing diagnostic audiology services in economically challenged regions. In: McPherson B, Brouillette R, eds. *Audiology in Developing Countries*. New York: Nova Science Publishers, Inc.; 2008:108-140.
18. Chonmaitree T, Revai K, Grady JJ, Clos A, Patel JA, Nair S, et al. Viral upper respiratory tract infection and otitis media complication in young children. *Clin Infect Dis* 2008; **46**(6): 815-823.
19. Fairchok MP, Martin ET, Chambers S, Kuypers J, Behrens M, Braun LE, et al. Epidemiology of viral respiratory tract infections in a prospective cohort of infants and toddlers attending daycare. *J Clin Virol* 2010; **49**(1): 16-20.
20. Hendley JO. Clinical virology of rhinoviruses. *Adv Virus Res* 1999; **54**: 453-466.
21. Nokso-Koivisto J, Hovi T, Pitkäranta A. Viral upper respiratory tract infections in young children with emphasis on acute otitis media. *Int J Pediatr Otorhinolaryngol* 2006; **70**(8): 1333-1342.
22. Pitkäranta A, Hayden FG. Rhinoviruses: important respiratory pathogens. *Ann Med* 1998; **30**(6): 529-537.
23. Ruohola A, Waris M, Allander T, Ziegler T, Heikkinen T, Ruuskanen O. Viral etiology of common cold in children, Finland. *Emerg Infect Dis* 2009; **15**(2): 344-346.
24. Abzug M. Nonpolio Enteroviruses. In: Kliegman R, Stanton B, St.Geme J, Schor N, Behrman R, eds. *Nelson Textbook of Pediatrics*. 19th ed. Philadelphia: Saunders; 2011: 1088-1094.
25. Hovi T, Hyypiä T. Enterovirukset, rinovirukset ja muut pikornavirukset. In: Huovinen P, Meri S, Peltola H, Vaara M, Vaheri A, Valtonen V, eds. *Mikrobiologia ja infektiosairaudet*. 1st ed. Jyväskylä: Duodecim; 2005: 450-462.
26. International Committee on Taxonomy of Viruses (ICTV). (Accessed 05/31, 2012, at <http://www.ictvonline.org/>).
27. Miller E, Williams J. Rhinoviruses. In: Kliegman R, Stanton B, St.Geme J, Schor N, Behrman R, eds. *Nelson Textbook of Pediatrics*. 19th ed. Philadelphia: Saunders; 2011: 1133-1134.
28. Greve JM, Davis G, Meyer AM, Forte CP, Yost SC, Marlor CW, et al. The major human rhinovirus receptor is ICAM-1. *Cell* 1989; **56**(5): 839-847.
29. Winther B, McCue K, Ashe K, Rubino JR, Hendley JO. Environmental contamination with rhinovirus and transfer to fingers of healthy individuals by daily life activity. *J Med Virol* 2007; **79**(10): 1606-1610.
30. Sattar SA, Karim YG, Springthorpe VS, Johnson-Lussenburg CM. Survival of human rhinovirus type 14 dried onto nonporous inanimate surfaces: effect of relative humidity and suspending medium. *Can J Microbiol* 1987; **33**(9): 802-806.



31. Johnston SL, Sanderson G, Pattemore PK, et al. Use of polymerase chain reaction for diagnosis of picornavirus infection in subjects with and without respiratory symptoms. *J Clin Microbiol* 1993;31:111-117.
32. Pappas DE, Hendley JO, Hayden FG, Winther B. Symptom profile of common colds in school-aged children. *Pediatr Infect Dis J* 2008; 27(1): 8-11.
33. Peltola V, Waris M, Osterback R, Susi P, Hyypia T, Ruuskanen O. Clinical effects of rhinovirus infections. *J Clin Virol* 2008; **43**(4): 411-414.
34. Monto AS, Fendrick AM, Sarnes MW. Respiratory illness caused by picornavirus infection: a review of clinical outcomes. *Clin Ther* 2001; **23**(10): 1615-1627.
35. Piotrowska Z, Vazquez M, Shapiro ED, Weibel C, Ferguson D, Landry ML, et al. Rhinoviruses are a major cause of wheezing and hospitalization in children less than 2 years of age. *Pediatr Infect Dis J* 2009; **28**(1): 25-29.
36. Smuts HE, Workman LJ, Zar HJ. Human rhinovirus infection in young African children with acute wheezing. *BMC Infect Dis* 2011; **11**: 65.
37. Fuji N, Suzuki A, Lupisan S, Sombrero L, Galang H, Kamigaki T, et al. Detection of human rhinovirus C viral genome in blood among children with severe respiratory infections in the Philippines. *PLoS One* 2011; **6**(11): e27247.
38. Khetsuriani N. Enterovirus surveillance--United States, 1970-2005. *MMWR Surveill Summ*. 2006; **55**(8): 1-20.
39. Pelkonen T, Roine I, Anjos E, Kaijalainen S, Roivainen M, Peltola H, et al. Picornaviruses in cerebrospinal fluid of children with meningitis in Luanda, Angola. *J Med Virol* 2012; **84**(7): 1080-1083.
40. Winther B, Hayden FG, Hendley JO. Picornavirus infections in children diagnosed by RT-PCR during longitudinal surveillance with weekly sampling: Association with symptomatic illness and effect of season. *J Med Virol* 2006; **78**(5): 644-650.
41. Nokso-Koivisto J, Rätty R, Blomqvist S, Kleemola M, Syrjänen R, Pitkäranta A, et al. Presence of specific viruses in the middle ear fluids and respiratory secretions of young children with acute Otitis Media. *J Med Virol* 2004; **72**(2): 241-248.
42. Vesa S, Kleemola M, Blomqvist S, Takala A, Kilpi T, Hovi T. Epidemiology of documented viral respiratory infections and acute otitis media in a cohort of children followed from two to twenty-four months of age. *Pediatr Infect Dis J* 2001; **20**(6):574-581.
43. Monto AS. Occurrence of respiratory virus: time, place and person. *Pediatr Infect Dis J* 2004; **23**(1 Suppl): S58-64.
44. Monto AS, Bryan ER, Ohmit S. Rhinovirus infections in Tecumseh, Michigan: frequency of illness and number of serotypes. *J Infect Dis* 1987; **156**(1): 43-49.
45. Du Prel JB, Puppe W, Grondahl B, Knuf M, Weigl JA, Schaaff F, et al. Are meteorological parameters associated with acute respiratory tract infections? *Clin Infect Dis* 2009; **49**(6): 861-868.
46. Karim YG, Ijaz MK, Sattar SA, Johnson-Lussenburg CM. Effect of relative humidity on the airborne survival of rhinovirus-14. *Can J Microbiol* 1985; **31**(11): 1058-1061.

47. Lina B, Valette M, Foray S, Luciani J, Stagnara J, See DM, et al. Surveillance of community-acquired viral infections due to respiratory viruses in Rhone-Alpes (France) during winter 1994 to 1995. *J Clin Microbiol* 1996; **34**(12): 3007-3011.
48. Onyango CO, Welch SR, Munywoki PK, Agoti CN, Bett A, Ngama M, et al. Molecular epidemiology of human rhinovirus infections in Kilifi, coastal Kenya. *J Med Virol* 2012; **84**(5): 823-831.
49. O'Callaghan-Gordo C, Díez-Padrisa N, Abacassamo F, Pérez-Breña P, Casas I, Alonso PL, et al. Viral acute respiratory infections among infants visited in a rural hospital of southern Mozambique. *Trop Med Int Health* 2011; **16**(9): 1054-1060.
50. Souza L, Ramos EA, Carvalho FM, Guedes VM, Souza LS, Rocha CM, et al. Viral respiratory infections in young children attending day care in urban Northeast Brazil. *Pediatr Pulmonol* 2003; **35**(3): 184-191.
51. Waitumbi JN, Kuypers J, Anyona SB, Koros JN, Polhemus ME, Gerlach J, et al. Outpatient upper respiratory tract viral infections in children with malaria symptoms in Western Kenya. *Am J Trop Med Hyg* 2010; **83**(5): 1010-1013.
52. Lin TY, Huang YC, Ning HC, Tsao KC. Surveillance of respiratory viral infections among pediatric outpatients in northern Taiwan. *J Clin Virol* 2004; **30**(1): 81-85.
53. O'Callaghan-Gordo C, Bassat Q, Morais L, Díez-Padrisa N, Machevo S, Nhampossa T, et al. Etiology and epidemiology of viral pneumonia among hospitalized children in rural Mozambique: A malaria endemic area with high prevalence of human immunodeficiency virus. *Pediatr Infect Dis J* 2011; **30**(1): 39-44.
54. Hazlett DTG, Bell TM, Tukei PM, Ademba GR, Ochieng WO, Magana JM, et al. Viral etiology and epidemiology of acute respiratory infections in children in Nairobi, Kenya. *Am J Trop Med Hyg* 1988; **39**(6): 632-640.
55. Yoshida LM, Suzuki M, Yamamoto T, Nguyen HA, Nguyen CD, Nguyen AT, et al. Viral pathogens associated with acute respiratory infections in central vietnamese children. *Pediatr Infect Dis J* 2010; **29**(1): 75-77.
56. Chew FT, Dorasingham S, Ling AE, Kumarasinghe G, Lee BW. Seasonal trends of viral respiratory tract infections in the tropics. *Epidemiol Infect* 1998; **121**(1): 121-128.
57. Pitkäranta A, Virolainen A, Jero J, Arruda E, Hayden FG. Detection of rhinovirus, respiratory syncytial virus, and coronavirus infections in acute otitis media by reverse transcriptase polymerase chain reaction. *Pediatrics* 1998; **102**(2 Pt 1):291-295.
58. Ruohola A, Meurman O, Nikkari S, Skottman T, Salmi A, Waris M, et al. Microbiology of acute otitis media in children with tympanostomy tubes: prevalences of bacteria and viruses. *Clin Infect Dis* 2006; **43**(11): 1417-1422.
59. Heikkinen T, Thint M, Chonmaitree T. Prevalence of various respiratory viruses in the middle ear during acute otitis media. *N Engl J Med* 1999; **340**(4): 260-264.
60. Verhoeff M, van der Veen EL, Rovers MM, Sanders EA, Schilder AG. Chronic suppurative otitis media: a review. *Int J Pediatr Otorhinolaryngol* 2006; **70**(1): 1-12.
61. Hamilton J. Chronic Otitis Media. In: Graham J, Scadding G, Bull P, eds. *Pediatric ENT*. 1st ed. Berlin Heidelberg: Springer-Verlag; 2007: 421-440.

62. Friedmann I. The pathology of otitis media. *J Clin Pathol* 1956; **9**(3): 229-236.
63. Bluestone CD. Epidemiology and pathogenesis of chronic suppurative otitis media: implications for prevention and treatment. *Int J Pediatr Otorhinolaryngol* 1998; **42**(3): 207-223.
64. Berman S. Otitis media in developing countries. *Pediatrics* 1995; **96**(1 Pt 1): 126-131.
65. Obi CL, Enweani IB, Giwa JO. Bacterial agents causing chronic suppurative otitis media. *East Afr Med J* 1995; **72**(6): 370-372.
66. Nwabuisi C, Ologe FE. Pathogenic agents of chronic suppurative otitis media in Ilorin, Nigeria. *East Afr Med J* 2002; **79**(4): 202-205.
67. Da Costa SS, Rosito LP, Dornelles C. Sensorineural hearing loss in patients with chronic otitis media. *Eur Arch Otorhinolaryngol* 2009; **266**(2): 221-224.
68. Couzos S, Lea T, Mueller R, Murray R, Culbong M. Effectiveness of ototopical antibiotics for chronic suppurative otitis media in Aboriginal children: a community-based, multicentre, double-blind randomised controlled trial. *Med J Aust* 2003; **179**(4): 185-190.
69. De Uzeda M, Rocha ER. Bacteriology of chronic otitis media affecting children living in Rio de Janeiro. *Ear Nose Throat J* 1989; **68**(6): 448, 450, 452, passim.
70. Van Hasselt P, van Kregten E. Treatment of chronic suppurative otitis media with ofloxacin in hydroxypropyl methylcellulose ear drops: a clinical/bacteriological study in a rural area of Malawi. *Int J Pediatr Otorhinolaryngol* 2002; **63**(1): 49-56.
71. Nyembue DT, Tshiswaka JM, Sabue MJ, Muyunga CK. Bacteriology of chronic suppurative otitis media in congolese children. *Acta Otorhinolaryngol Belg* 2003; **57**(3): 205-208.
72. Eason RJ, Harding E, Nicholson R, Nicholson D, Pada J, Gathercole J. Chronic suppurative otitis media in the Solomon Islands: a prospective, microbiological, audiometric and therapeutic survey. *N Z Med J* 1986; **99**(812): 812-5.
73. Hussain MA, Ali EM, Ahmed HS. Otitis media in Sudanese children: presentation and bacteriology. *East Afr Med J* 1991; **68**(9): 679-685.
74. Morris PS, Leach AJ, Silberberg P, Mellon G, Wilson C, Hamilton E, et al. Otitis media in young Aboriginal children from remote communities in Northern and Central Australia: a cross-sectional survey. *BMC Pediatr* 2005; **5**: 27.
75. Bruneau S, Ayukawa H, Proulx JF, Baxter JD, Kost K. Longitudinal observations (1987-1997) on the prevalence of middle ear disease and associated risk factors among Inuit children of Inukjuak, Nunavik, Quebec, Canada. *Int J Circumpolar Health* 2001; **60**(4): 632-639.
76. Baxter JD. Otitis media in Inuit children in the Eastern Canadian Arctic--an overview--1968 to date. *Int J Pediatr Otorhinolaryngol* 1999; **49 Suppl 1**: S165-168.
77. Homoe P. Otitis media in Greenland. Studies on historical, epidemiological, microbiological, and immunological aspects. *Int J Circumpolar Health* 2001; **60 Suppl 2**: 1-54.
78. Leach AJ. Otitis media in Australian Aboriginal children: an overview. *Int J Pediatr Otorhinolaryngol* 1999; **49 Suppl 1**: S173-178.

79. Bastos I, Reimer A, Lundgren K. Chronic otitis media and hearing loss in urban schoolchildren in Angola - A prevalence study. *Journal of Audiological Medicine* 1993; **2**: 129-140.
80. Hatcher J, Smith A, Mackenzie I, Thompson S, Bal I, Macharia I, et al. A prevalence study of ear problems in school children in Kiambu district, Kenya, May 1992. *Int J Pediatr Otorhinolaryngol* 1995; **33**(3): 197-205.
81. Lasisi AO, Sulaiman OA, Afolabi OA. Socio-economic status and hearing loss in chronic suppurative otitis media in Nigeria. *Ann Trop Paediatr* 2007; **27**(4): 291-296.
82. Olatoke F, Ologe FE, Nwawolo CC, Saka MJ. The prevalence of hearing loss among schoolchildren with chronic suppurative otitis media in Nigeria, and its effect on academic performance. *Ear Nose Throat J* 2008; **87**(12): E19.
83. Causes of death 2008, Summary tables. May 2011. World Health Organization, Geneva. (Accessed 06/10, 2012, at <http://www.who.int/evidence/bod>).
84. Monasta L, Ronfani L, Marchetti F, Montico M, Vecchi Brumatti L, Bavcar A, et al. Burden of disease caused by otitis media: systematic review and global estimates. *PLoS One* 2012; **7**(4): e36226.
85. Fliss DM, Shoham I, Leiberman A, Dagan R. Chronic suppurative otitis media without cholesteatoma in children in southern Israel: incidence and risk factors. *Pediatr Infect Dis J* 1991; **10**(12): 895-899.
86. Elemraïd MA, Mackenzie IJ, Fraser WD, Brabin BJ. Nutritional factors in the pathogenesis of ear disease in children: A systematic review. *Ann Trop Paediatr* 2009; **29**(2): 85-99.
87. Lasisi AO. The role of retinol in the etiology and outcome of suppurative otitis media. *Eur Arch Otorhinolaryngol* 2009; **266**(5): 647-652.
88. Schmitz J, West KP Jr, Khatri SK, Wu L, Leclerc SC, Karna SL, et al. Vitamin A supplementation in preschool children and risk of hearing loss as adolescents and young adults in rural Nepal: randomised trial cohort follow-up study. *BMJ* 2012; **344**: d7962.
89. Macfadyen CA, Acuin JM, Gamble C. Topical antibiotics without steroids for chronically discharging ears with underlying eardrum perforations. *Cochrane Database Syst Rev* 2005; **4**: CD004618.
90. Dohar JE, Garner ET, Nielsen RW, Biel MA, Seidlin M. Topical ofloxacin treatment of otorrhea in children with tympanostomy tubes. *Arch Otolaryngol Head Neck Surg* 1999; **125**(5): 537-545.
91. Smith AW, Hatcher J, Mackenzie IJ, Thompson S, Bal I, Macharia I, et al. Randomised controlled trial of treatment of chronic suppurative otitis media in Kenyan schoolchildren. *Lancet* 1996; **348**(9035): 1128-1133.
92. Hannley MT, Denny JC 3rd, Holzer SS. Use of ototopical antibiotics in treating 3 common ear diseases. *Otolaryngol Head Neck Surg* 2000; **122**(6): 934-940.
93. Tong MC, Woo JK, van Hasselt CA. A double-blind comparative study of ofloxacin otic drops versus neomycin-polymyxin B-hydrocortisone otic drops in the medical treatment of chronic suppurative otitis media. *J Laryngol Otol* 1996; **110**(4): 309-314.

94. Macfadyen C, Gamble C, Garner P, Macharia I, Mackenzie I, Mugwe P, et al. Topical quinolone vs. antiseptic for treating chronic suppurative otitis media: a randomized controlled trial. *Clin Otolaryngol* 2005; **30**(2): 193-194.
95. Minja BM, Moshi NH, Ingvarsson L, Bastos I, Grenner J. Chronic suppurative otitis media in Tanzanian school children and its effects on hearing. *East Afr Med J* 2006; **83**(6): 322-325.
96. Olusesi AD. Otitis media as a cause of significant hearing loss among Nigerians. *Int J Pediatr Otorhinolaryngol* 2008; **72**(6): 787-792.
97. Van der Veen EL, Schilder AG, van Heerbeek N, Verhoeff M, Zielhuis GA, Rovers MM. Predictors of chronic suppurative otitis media in children. *Arch Otolaryngol Head Neck Surg* 2006; **132**(10): 1115-1118.
98. Brook I. The role of anaerobic bacteria in chronic suppurative otitis media in children: implications for medical therapy. *Anaerobe* 2008; **14**(6): 297-300.
99. Brook I, Santosa G. Microbiology of chronic suppurative otitis media in children in Surabaya, Indonesia. *Int J Pediatr Otorhinolaryngol* 1995; **31**(1): 23-28.
100. Kong K, Coates HL. Natural history, definitions, risk factors and burden of otitis media. *Med J Aust* 2009; **191**(9 Suppl): S39-43.
101. Post JC, Hiller NL, Nistico L, Stoodley P, Ehrlich GD. The role of biofilms in otolaryngologic infections: update 2007. *Curr Opin Otolaryngol Head Neck Surg* 2007; **15**(5): 347-351.
102. Wu JF, Jin Z, Yang JM, Liu YH, Duan ML. Extracranial and intracranial complications of otitis media: 22-year clinical experience and analysis. *Acta Otolaryngol* 2012; **132**(3): 261-265.
103. Matanda RN, Muyunga KC, Sabue MJ, Creten W, Van de Heyning P. Chronic suppurative otitis media and related complications at the University Clinic of Kinshasa. *B-ENT* 2005; **1**(2): 57-62.
104. Kärjä J, Jokinen K, Seppälä A. Destruction of ossicles in chronic otitis media. *J Laryngol Otol* 1976; **90**(6): 509-518.
105. Tos M. Pathology of the ossicular chain in various chronic middle ear diseases. *J Laryngol Otol* 1979; **93**(8): 769-780.
106. Jeng FC, Tsai MH, Brown CJ. Relationship of preoperative findings and ossicular discontinuity in chronic otitis media. *Otol Neurotol* 2003; **24**(1): 29-32.
107. El-Sayed Y. Bone conduction impairment in uncomplicated chronic suppurative otitis media. *Am J Otolaryngol* 1998; **19**(3): 149-153.
108. Papp Z, Rezes S, Jókay I, Sziklai I. Sensorineural hearing loss in chronic otitis media. *Otol Neurotol* 2003; **24**(2): 141-144.
109. Engel F, Blatz R, Schliebs R, Palmer M, Bhakdi S. Bacterial cytolysin perturbs round window membrane permeability barrier in vivo: possible cause of sensorineural hearing loss in acute otitis media. *Infect Immun* 1998; **66**(1): 343-346.
110. Matz G, Rybak L, Roland PS, Hannley M, Friedman R, Manolidis S. Ototoxicity of ototopical antibiotic drops in humans. *Otolaryngol Head Neck Surg* 2004; **130**(3 Suppl): S79-82.

111. Akinpelu OV, Amusa YB, Komolafe EO, Adeolu AA, Oladele AO, Ameye SA. Challenges in management of chronic suppurative otitis media in a developing country. *J Laryngol Otol* 2008; **122**(1): 16-20.
112. Okafor BC. The chronic discharging ear in Nigeria. *J Laryngol Otol* 1984; **98**(2): 113-119.
113. Kangsanarak J. Extracranial and intracranial complications of suppurative otitis media. Report of 102 cases. *J Laryngol Otol* 1993; **107**(11): 999-1004.
114. Seely DR, Gloyd SS, Wright AD, Norton SJ. Hearing loss prevalence and risk factors among Sierra Leonean children. *Arch Otolaryngol Head Neck Surg* 1995; **121**(8): 853-858.
115. Lasisi AO, Olaniyan FA, Muibi SA, Azeez IA, Abdulwasiiu KG, Lasisi TJ, et al. Clinical and demographic risk factors associated with chronic suppurative otitis media. *Int J Pediatr Otorhinolaryngol* 2007; **71**(10): 1549-1554.
116. Lasisi AO, Olayemi O, Irabor AE. Early onset otitis media: risk factors and effects on the outcome of chronic suppurative otitis media. *Eur Arch Otorhinolaryngol* 2008; **265**(7): 765-768.
117. Kaplan GJ, Fleshman JK, Baum C, Clark PS. Long-term effects of otitis media: a ten-year cohort study of Alaskan Eskimo children. *Pediatrics* 1973; **52**(4): 577-585.
118. Ebrahim SH, Khoja TA, Elachola H, Atrash HK, Memish Z, Johnson A. Children who come and go: the state of sickle cell disease in resource-poor countries. *Am J Prev Med* 2010; **38**(4 Suppl): S568-570.
119. Weatherall DJ. The inherited diseases of hemoglobin are an emerging global health burden. *Blood* 2010; **115**(22): 4331-4336.
120. Modell B, Darlison M. Global epidemiology of haemoglobin disorders and derived service indicators. *Bull World Health Organ* 2008; **86**(6): 480-487.
121. Williams TN, Mwangi TW, Wambua TS, Alexander ND, Kortok M, Snow RW, et al. Sickle cell trait and the risk of *Plasmodium falciparum* malaria and other childhood diseases. *J Infect Dis* 2005; **192**(1): 178-186.
122. Grosse SD, Odam E, Atrash HK, Amendah DD, Piel FB, Williams TN. Sickle cell disease in Africa: a neglected cause of early childhood mortality. *Am J Prev Med* 2011; **41**(6 Suppl 4): S398-405.
123. Ramakrishnan M, Moisi JC, Klugman KP, Iglesias JM, Grant LR, Mpoudi-Etame M, et al. Increased risk of invasive bacterial infections in African people with sickle-cell disease: a systematic review and meta-analysis. *Lancet Infect Dis* 2010; **10**(5): 329-337.
124. Rees DC, Williams TN, Gladwin MT. Sickle-cell disease. *Lancet* 2010; **376**(9757): 2018-2031.
125. Sickle cell disease prevention and control. World Health Organization: Regional Office for Africa. (Accessed 05/09, 2012, at <http://www.afro.who.int/en/>).
126. Serjeant GR, Loy LL, Crowther M, Hambleton IR, Thame M. Outcome of pregnancy in homozygous sickle cell disease. *Obstet Gynecol* 2004; **103**(6): 1278-1285.
127. eBaun MR. Sickle Cell Disease. In: Kliegman R, Stanton B, St.Geme J, Schor N, Behrman R, eds. *Nelson Textbook of Pediatrics*. 19th ed. Philadelphia: Elsevier Saunders; 2011: 1663-1670.

128. Di Sabatino A, Carsetti R, Corazza GR. Post-splenectomy and hyposplenic states. *Lancet* 2011; **378**(9785): 86-97.
129. Van-Dunem JC, Alves JG, Bernardino L, Figueiroa JN, Braga C, do Nascimento Mde L, et al. Factors associated with sickle cell disease mortality among hospitalized Angolan children and adolescents. *West Afr J Med* 2007; **26**(4): 269-273.
130. Adamkiewicz T. Invasive infections and sickle-cell disease. *Lancet Infect Dis* 2010; **10**(9): 593; author reply 593-4.
131. Knight-Madden J, Serjeant GR. Invasive pneumococcal disease in homozygous sickle cell disease: Jamaican experience 1973-1997. *J Pediatr* 2001; **138**(1): 65-70.
132. De Montalembert M, Brousse V, Zahar JR. Pneumococcal prophylaxis for children with sickle cell disease in Africa. *Arch Dis Child* 2008; **93**(8): 715-716.
133. Makani J, Cox SE, Soka D, Komba AN, Oruo J, Mwantemi H, et al. Mortality in sickle cell anemia in Africa: a prospective cohort study in Tanzania. *PLoS One* 2011; **6**(2): e14699.
134. Cox SE, Makani J, Fulford AJ, Komba AN, Soka D, Williams TN, et al. Nutritional status, hospitalization and mortality among patients with sickle cell anemia in Tanzania. *Haematologica* 2011; **96**(7): 948-953.
135. Weatherall MW, Higgs DR, Weiss H, Weatherall DJ, Serjeant GR. Phenotype/genotype relationships in sickle cell disease: a pilot twin study. *Clin Lab Haematol* 2005; **27**(6): 384-390.
136. De Montalembert M. Management of sickle cell disease. *BMJ* 2008; **337**(7670): 626-630.
137. Ryan K, Bain BJ, Worthington D, James J, Plews D, Mason A, et al. Significant haemoglobinopathies: guidelines for screening and diagnosis. *Br J Haematol* 2010; **149**(1): 35-49.
138. Vichinsky E, Hurst D, Earles A, Kleman K, Lubin B. Newborn screening for sickle cell disease: effect on mortality. *Pediatrics* 1988; **81**(6): 749-755.
139. Alabi S, Ernest K, Eletta P, Owolabi A, Afolabi A, Suleiman O. Otological findings among Nigerian children with sickle cell anaemia. *Int J Pediatr Otorhinolaryngol* 2008; **72**(5): 659-663.
140. De Castro Silva IM, Magalhaes IQ, Toscano RA, Gandolfi L, Pratesi R. Auditory-evoked response analysis in Brazilian patients with sickle cell disease. *Int J Audiol* 2010; **49**(4): 272-276.
141. Elwany S, Kamel T. Sensorineural hearing loss in sickle cell crisis. *Laryngoscope* 1988; **98**(4): 386-389.
142. Forman-Franco B, Karayalcin G, Mandel DD, Abramson AL. The evaluation of auditory function in homozygous sickle cell disease. *Otolaryngol Head Neck Surg* 1982; **90**(6): 850-856.
143. Mgbor N, Emodi I. Sensorineural hearing loss in Nigerian children with sickle cell disease. *Int J Pediatr Otorhinolaryngol* 2004; **68**(11): 1413-1416.
144. Odetoyinbo O, Adekile A. Sensorineural hearing loss in children with sickle cell anemia. *Ann Otol Rhinol Laryngol* 1987; **96**(3 Pt 1): 258-260.



## References

145. Tsibulevskaya G, Oburra H, Aluoch JR. Sensorineural hearing loss in patients with sickle cell anaemia in Kenya. *East Afr Med J* 1996; **73**(7): 471-473.
146. Ajulo SO, Osiname AI, Myatt HM. Sensorineural hearing loss in sickle cell anaemia--a United Kingdom study. *J Laryngol Otol* 1993; **107**(9): 790-794.
147. Todd GB, Serjeant GR, Larson MR . Sensori-neural hearing loss in Jamaicans with SS disease. *Acta Otolaryngol* 1973; **76**(4): 268-272.
148. Samperi P, Bertuna G, Rossi G, Poli G, Serra A. Sensorineural hearing loss in sickle cell disease patients in Sicily. *Minerva Pediatr* 2005; **57**(5): 285-288.
149. Atsina KK, Ankra-Badu G. Sensorineural hearing loss in Ghanaiaians with sickle cell anaemia. *Trop Geogr Med* 1988; **40**(3): 205-208.
150. Aderibigbe A, Ologe FE, Oyejola BA. Hearing thresholds in sickle cell anemia patients: emerging new trends? *J Natl Med Assoc* 2005; **97**(8): 1135-1142.
151. Jovanovic-Bateman L, Hedreville R. Sensorineural hearing loss with brainstem auditory evoked responses changes in homozygote and heterozygote sickle cell patients in Guadeloupe (France). *J Laryngol Otol* 2006; **120**(8): 627-630.
152. Piltcher O, Cigana L, Friedrich J, Ribeiro FA, da Costa SS. Sensorineural hearing loss among sickle cell disease patients from southern Brazil. *Am J Otolaryngol* 2000; **21**(2): 75-79.
153. Morgenstein KM, Manace ED. Temporal bone histopathology in sickle cell disease. *Laryngoscope* 1969; **79**(12): 2172-2180.
154. Hotaling AJ, Hillstrom RP, Bazell C. Sickle cell crisis and sensorineural hearing loss: case report and discussion. *Int J Pediatr Otorhinolaryngol* 1989; **17**(3): 207-211.
155. Belal A Jr. The effects of vascular occlusion on the human inner ear. *Journal of Laryngology & Otology* 1979; **93**(10): 955-968.
156. Liu BP, Saito N, Wang JJ, Mian AZ, Sakai O. Labyrinthitis ossificans in a child with sickle cell disease: CT and MRI findings. *Pediatr Radiol* 2009; **39**(9): 999-1001.
157. Gentry B, Davis P, Dancer J. Failure rates of young patients with sickle cell disease on a hearing screening test. *Percept Mot Skills* 1997; **84**(2): 434.
158. oussi A, Zafeiriou DI, Kontzoglou G, Tsatra I, Noussios G, Athanassiou M. Hearing loss in children with sickle cell disease. *Acta Otorhinolaryngol Belg* 2001; **55**(3): 235-239.
159. MacDonald CB, Bauer PW, Cox LC, McMahon L. Otologic findings in a pediatric cohort with sickle cell disease. *Int J Pediatr Otorhinolaryngol* 1999; **47**(1): 23-28.
160. Al Okbi MH, Alkindi S, Al Abri RK, Mathew J, Nagwa AA, Pathare AV. Sensorineural hearing loss in sickle cell disease--a prospective study from Oman. *Laryngoscope* 2011; **121**(2): 392-396.
161. Elola A, Hien EM, Ouattara M, FataoFataho B, Kouassi B. Drépanocytose majeure et hypoacousie : À propos de 112 cas relevés à Yopougon, Côte d'Ivoire. *Bull Soc Pathol Exot* 2009; **102**(3): 173-174.
162. Serjeant GR, Norman W, Todd GB. The internal auditory canal and sensori-neural hearing loss in homozygous sickle cell disease. *J Laryngol Otol* 1975; **89**(4): 453-455.



163. Williams TN, Uyoga S, Macharia A, Ndila C, McAuley CF, Opi DH, et al. Bacteraemia in Kenyan children with sickle-cell anaemia: a retrospective cohort and case-control study. *Lancet* 2009; **374**(9698): 1364-1370.
164. Lajeunesse M. The immunocompromised child. In: Graham J, Scadding G, Bull P, eds. *Pediatric ENT*. 1st ed. Berlin Heidelberg: Springer-Verlag; 2007: 83-93.
165. Wilson NW, Hogan MB. Otitis media as a presenting complaint in childhood immunodeficiency diseases. *Curr Allergy Asthma Rep* 2008; **8**(6): 519-524.
166. Donegan JO, Lobel JS, Gluckman JL. Otolaryngologic manifestations of sickle cell disease. *Am J Otolaryngol* 1982; **3**(2): 141-144.
167. Alkan O, Kizilkilic E, Kizilkilic O, Yildirim T, Karaca S, Yeral M, et al. Cranial involvement in sickle cell disease. *Eur J Radiol* 2010; **76**(2): 151-156.
168. Alli NA, Wainwright RD, Mackinnon D, Poyiadjis S, Naidu G. Skull bone infarctive crisis and deep vein thrombosis in homozygous sickle cell disease- case report and review of the literature. *Hematology* 2007; **12**(2): 169-174.
169. Saito N, Nadgir RN, Flower EN, Sakai O. Clinical and radiologic manifestations of sickle cell disease in the head and neck. *Radiographics* 2010; **30**(4): 1021-1034.
170. Abou-Elhamd K. Otorhinolaryngological manifestations of sickle cell disease. *Int J Pediatr Otorhinolaryngol* 2012; **76**(1): 1-4.
171. Samuels MP, Stebbens VA, Davies SC, Picton-Jones E, Southall DP. Sleep related upper airway obstruction and hypoxaemia in sickle cell disease. *Arch Dis Child* 1992; **67**(7): 925-929.
172. Yogev R, Chadwick EG. Acquired Immunodeficiency Syndrome (Human Immunodeficiency Virus). In: Kliegman R, Stanton B, St. Geme J, Schor N, Behrman R, eds. *Nelson textbook of pediatrics*. 19th ed. Philadelphia: Elsevier Saunders; 2011: 1157-1177.
173. Spira R, Lepage P, Msellati P, Van De Perre P, Leroy V, Simonon A, et al. Natural history of human immunodeficiency virus type 1 infection in children: a five-year prospective study in Rwanda. Mother-to-Child HIV-1 Transmission Study Group. *Pediatrics* 1999; **104**(5): e56.
174. Newell ML, Coovadia H, Cortina-Borja M, Rollins N, Gaillard P, Dabis F. Mortality of infected and uninfected infants born to HIV-infected mothers in Africa: a pooled analysis. *Lancet* 2004; **364**(9441): 1236-1243.
175. Grant AD, De Cock KM. ABC of AIDS. HIV infection and AIDS in the developing world. *BMJ* 2001; **322**(7300): 1475-1478.
176. UNAIDS report on the global AIDS epidemic 2010. World Health Organization. (Accessed 05/10, 2010, at <http://issuu.com/unaids/docs/>).
177. World Health Organization. WHO case definitions of HIV for surveillance and revised clinical staging and immunological classification of HIV-related disease in adults and children. Geneva; 2007.
178. World Health Organization. Antiretroviral therapy of HIV infection in infants and children: towards universal access: recommendations for a public health approach - 2010 revision. Geneva; 2010.

179. Chi BH, Limbada MI, Giganti MJ, Li MS, Bweupe M, Musonda P, et al. Non-virologic algorithms for predicting HIV infection among HIV-exposed infants under 12 weeks of age. *Pediatr Infect Dis J* 2013; **32**(2): 151-156.
180. Nielsen-Saines K, Watts DH, Veloso VG, Bryson YJ, Joao EC, Pilotto JH, et al. Three Postpartum Antiretroviral Regimens to Prevent Intrapartum HIV Infection. *N Engl J Med* 2012; **366**(25): 2368-2379.
181. UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance. Epidemiological Fact Sheets on HIV and AIDS. World Health Organization. (Accessed 05/10, 2010, at <http://unaids.org/en/resources/presscentre/factsheets/>)
182. Chow JH, Stern JC, Kaul A, Pincus RL, Gromisch DS. Head and neck manifestations of the acquired immunodeficiency syndrome in children. *Ear Nose Throat J* 1990; **69**(6): 416-419, 422-423.
183. De Vincentiis GC, Sitzia E, Bottero S, Giuzio L, Simonetti A, Rossi P. Otolaryngologic manifestations of pediatric immunodeficiency. *Int J Pediatr Otorhinolaryngol* 2009; **73 Suppl 1**: S42-48.
184. Singh A, Georgalas C, Patel N, Papesch M. ENT presentations in children with HIV infection. *Clin Otolaryngol Allied Sci* 2003; **28**(3): 240-243.
185. Hadfield PJ, Birchall MA, Novelli V, Bailey CM. The ENT manifestations of HIV infection in children. *Clin Otolaryngol Allied Sci* 1996; **21**(1): 30-36.
186. haloryoo S, Chotpitayasunondh T, Chiengmai PN. AIDS in ENT in children. *Int J Pediatr Otorhinolaryngol* 1998; **44**(2): 103-107.
187. Williams MA. Head and neck findings in pediatric acquired immune deficiency syndrome. *Laryngoscope* 1987; **97**(6): 713-716.
188. Gondim LA, Zonta RF, Fortkamp E, Schmeling RO. Otorhinolaryngological manifestations in children with human immunodeficiency virus infection. *Int J Pediatr Otorhinolaryngol* 2000; **54**(2-3): 97-102.
189. Prasad HK, Bhojwani KM, Shenoy V, Prasad SC. HIV manifestations in otolaryngology. *Am J Otolaryngol* 2006; **27**(39): 179-185.
190. Benito MB, Sampelayo TH, Gurbindo D, Sanchez-Ramon BS, Gomez EM, Munoz-Fernandez MA. Adenoidal tissue mass as a clinical guide of disease evolution in vertically HIV-1 infected children. *Int J Pediatr Otorhinolaryngol* 1999; **51**(3): 145-155.
191. Chao CK, Czechowicz JA, Messner AH, Alarcón J, Kolevic Roca L, Larragán Rodriguez MM, et al. High prevalence of hearing impairment in HIV-infected Peruvian children. *Otolaryngol Head Neck Surg* 2012; **146**(2): 259-265.
192. Palacios GC, Montalvo MS, Fraire MI, Leon E, Alvarez MT, Solorzano F. Audiologic and vestibular findings in a sample of human immunodeficiency virus type-1-infected Mexican children under highly active antiretroviral therapy. *Int J Pediatr Otorhinolaryngol* 2008; **72**(11): 1671-1681.
193. Matas CG, Santos Filha VA, Juan KR, Pinto FR, Goncalves IC. Audiological manifestations in children and adults with AIDS. *Pro Fono* 2010; **22**(3): 269-274.

194. Matas CG, Leite RA, Magliaro FC, Goncalves IC. Audiological and electrophysiological evaluation of children with acquired immunodeficiency syndrome (AIDS). *Braz J Infect Dis* 2006; **10**(4): 264-268.
195. Khoza-Shangase K. Highly active antiretroviral therapy: Does it Sound toxic? *J Pharm Bioallied Sci* 2011; **3**(1): 142-153.
196. Prazic M, Salaj B. Ototoxicity with children caused by streptomycin. *Audiology* 1975; **14**(2):173-176.
197. Harris T, Peer S, Fagan JJ. Audiological monitoring for ototoxic tuberculosis, human immunodeficiency virus and cancer therapies in a developing world setting. *J Laryngol Otol* 2012; **126**(6): 548-551.
198. Saunders JE, Greinwald JH, Vaz S, Guo Y. Aminoglycoside ototoxicity in Nicaraguan children: patient risk factors and mitochondrial DNA results. *Otolaryngol Head Neck Surg* 2009; **140**(1): 103-107.
199. Prada JL, Torre-Cisneros J, Kindelan JM, Jurado J, Villanueva JL, Navarro M, et al. Deafness and blindness in a HIV-positive patient with cryptococcal meningitis. *Postgrad Med J* 1996; **72**(851): 575.
200. Michaels L, Soucek S, Liang J. The ear in the acquired immunodeficiency syndrome: I. Temporal bone histopathologic study. *Am J Otol* 1994; **15**(4): 515-522.
201. Shapiro NL, Novelli V. Otitis media in children with vertically-acquired HIV infection: the Great Ormond Street Hospital experience. *Int J Pediatr Otorhinolaryngol* 1998; **45**(1): 69-75.
202. Chen AY, Ohlms LA, Stewart MG, Kline MW. Otolaryngologic disease progression in children with human immunodeficiency virus infection. *Arch Otolaryngol Head Neck Surg* 1996; **122**(12): 1360-1363.
203. Barnett ED, Klein JO, Pelton SI, Luginbuhl LM. Otitis media in children born to human immunodeficiency virus-infected mothers. *Pediatr Infect Dis J* 1992; **11**(5): 360-364.
204. Karpakis J, Rabie H, Howard J, Janse van Rensburg A, Cotton MF. Otorrhoea is a marker for symptomatic disease in HIV-infected children. *S Afr Med J* 2007; **97**(12): 1292-1294.
205. Bagenda D, Nassali A, Kalyesubula I, Sherman B, Drotar D, Boivin MJ, et al. Health, neurologic, and cognitive status of HIV-infected, long-surviving, and antiretroviral-naïve Ugandan children. *Pediatrics* 2006; **117**(3): 729-740.
206. Nkrumah FK, Choto RG, Emmanuel J, Kumar R. Clinical presentation of symptomatic human immuno-deficiency virus in children. *Cent Afr J Med* 1990; **36**(5): 116-120.
207. Weber R, Pinheiro Neto CD, Miziara ID, Araujo Filho BC. HAART impact on prevalence of chronic otitis media in Brazilian HIV-infected children. *Braz J Otorhinolaryngol* 2006; **72**(4): 509-514.
208. Okechukwu AA, Gambo D, Okechukwu OI. The clinical features of paediatric HIV/AIDS at presentation at the University of Abuja Teaching Hospital, Gwagwalada. *Niger J Med* 2008; **17**(4): 433-438.

209. Ogunbosi BO, Oladokun RE, Brown BJ, Osinusi KI. Prevalence and clinical pattern of paediatric HIV infection at the University College Hospital, Ibadan, Nigeria: a prospective cross-sectional study. *Ital J Pediatr* 2011; **37**: 29.
210. Ugochukwu EF, Ezechukwu CC, Undie N, Akujobi C. Pattern of pathogens in ear discharge of HIV-infected children in Nnewi, Southeast Nigeria. *Niger J Clin Pract* 2007; **10**(2): 130-136.
211. Lu X, Holloway B, Dare RK, Kuypers J, Yagi S, Williams JV, et al. Real-time reverse transcription-PCR assay for comprehensive detection of human rhinoviruses. *J Clin Microbiol* 2008; **46**(2): 533-539.
212. Kilpatrick DR, Yang CF, Ching K, Vincent A, Iber J, Campagnoli R, et al. Rapid group-, serotype-, and vaccine strain-specific identification of poliovirus isolates by real-time reverse transcription-PCR using degenerate primers and probes containing deoxyinosine residues. *J Clin Microbiol* 2009; **47**(6): 1939-1941.
213. Mulholland EK, Simoes EA, Costales MO, McGrath EJ, Manalac EM, Gove S. Standardized diagnosis of pneumonia in developing countries. *Pediatr Infect Dis J* 1992; **11**(2): 77-81.
214. Harari M, Shann F, Spooner V, Meisner S, Carney M, de Campo J. Clinical signs of pneumonia in children. *Lancet* 1991; **338**(8772): 928-930.
215. Hearing impairment grades. World Health Organization: Prevention of Blindness and Deafness. (Accessed 06/14, 2012, at <http://www.who.int/pbd/deafness/en/>).
216. Dr. Luis Bernardino: Personal communication; Luanda 2007.
217. Schlaudecker EP, Heck JP, Macintyre ET, Martinez R, Dodd CN, McNeal MM, et al. Etiology and seasonality of viral respiratory infections in rural Honduran children. *Pediatr Infect Dis J* 2012; **31**(11): 1113-1118.
218. Niang MN, Diop OM, Sarr FD, Goudiaby D, Malou-Sompy H, Ndiaye K, et al. Viral etiology of respiratory infections in children under 5 years old living in tropical rural areas of Senegal: The EVIRA project. *J Med Virol* 2010; **82**(5): 866-872.
219. Peltola V, Ruuskanen O. Editorial commentary: Respiratory viral infections in developing countries: common, severe, and unrecognized. *Clin Infect Dis* 2008; **46**(1): 58-60.
220. Garcia-Garcia ML, Calvo C, Pozo F, Villadangos PA, Perez-Brena P, Casas I. Spectrum of Respiratory Viruses in Children with Community Acquired Pneumonia. *Pediatr Infect Dis J* 2012; **31**(8): 808-813.
221. Fry AM, Lu X, Olsen SJ, Chittaganpitch M, Sawatwong P, Chantira S, et al. Human rhinovirus infections in rural Thailand: epidemiological evidence for rhinovirus as both pathogen and bystander. *PLoS One* 2011; **6**(3): e17780.
222. Venter M, Lassaunière M, Kresfelder TR, Westerberg Y, Visser A. Contribution of common and recently described respiratory viruses to annual hospitalizations in children in South Africa. *J Med Virol* 2011; **83**(8): 1458-1468.
223. Peltola V, Heikkinen T, Ruuskanen O, Jartti T, Hovi T, Kilpi T, et al. Temporal association between rhinovirus circulation in the community and invasive pneumococcal disease in children. *Pediatr Infect Dis J* 2011; **30**(6): 456-461.

224. Launes C, de-Sevilla MF, Selva L, Garcia-Garcia JJ., Pallares R, Muños-Almagro C. Viral coinfection in children less than five years old with invasive pneumococcal disease. *Pediatr Infect Dis J* 2012; **31**(6): 650-653.
225. Bastos I, Janzon L, Lundgren K, Reimer A. Otitis media and hearing loss in children attending an ENT clinic in Luanda, Angola. *Int J Pediatr Otorhinolaryngol* 1990; **20**(2): 137-148.
226. Raju KG, Unnykrishnan P, Nayar RJ, Dutt S, Macaden R. Reliability of conventional ear swabs in tubotympanic CSOM. *J Laryngol Otol* 1990; **104**(6): 460-462.
227. Odetoyinbo O. Early diagnosis of tuberculous otitis media. *J Laryngol Otol* 1988; **102**(2): 133-135.
228. Meher R, Singh I, Yadav SP, Gathwala G. Tubercular otitis media in children. *Otolaryngol Head Neck Surg* 2006; **135**(4): 650-652.
229. Tran KD, Koprowska IA, Rao S, Sundaram R, Goldstein NA. Myringotomy and tympanostomy tube placement in children with sickle cell disease. *Int J Pediatr Otorhinolaryngol* 2004; **68**(10): 1301-1305.
230. Aghamohammadi A, Moazzami K, Rezaei N, Karimi A, Movahedi M, Gharagozlou M, et al. ENT manifestations in Iranian patients with primary antibody deficiencies. *J Laryngol Otol* 2008; **122**(4): 409-413.
231. Urschel S. Otitis media in children with congenital immunodeficiencies. *Curr Allergy Asthma Rep* 2010; **10**(6): 425-433.
232. Serjeant GR. Mortality from sickle cell disease in Africa. *BMJ* 2005; **330**(7489): 432-433.
233. Kizito ME, Mworozi E, Ndugwa C, Serjeant GR. Bacteraemia in homozygous sickle cell disease in Africa: is pneumococcal prophylaxis justified? *Arch Dis Child* 2007; **92**(1): 21-23.
234. Thomas EM. Recent trends in upper respiratory infections, ear infections and asthma among young Canadian children. *Health Rep* 2010; **21**(4): 47-52.
235. Paradise JL, Rockette HE, Colborn DK, Bernard BS, Smith CG, Kurs-Lasky M, et al. Otitis media in 2253 Pittsburgh-area infants: Prevalence and risk factors during the first two years of life. *Pediatrics* 1997; **99**(3): 318-333.
236. Olusanya BO, Okolo AA, Aderemi AA. Predictors of hearing loss in school entrants in a developing country. *J Postgrad Med* 2004; **50**(3): 173-179.
237. Olusanya BO, Okolo AA, Ijaduola GT. The hearing profile of Nigerian school children. *Int J Pediatr Otorhinolaryngol* 2000; **55**(3): 173-179.
238. Al Khabori M, Kumar S, Khandekar R. Magnitude of impacted earwax in Oman, its impact on hearing impairment and economic burden of earwax on health services. *Indian J Med Sci* 2007; **61**(5): 278-285.
239. Czechowicz JA, Messner AH, Alarcon-Matutti E, Alarcon J, Quinones-Calderon G, Montano S, et al. Hearing impairment and poverty: the epidemiology of ear disease in Peruvian schoolchildren. *Otolaryngol Head Neck Surg* 2010; **142**(2): 272-277.

240. Memel D, Langley C, Watkins C, Laue B, Birchall M, Bachmann M. Effectiveness of ear syringing in general practice: a randomised controlled trial and patients' experiences. *Br J Gen Pract* 2002; **52**(484): 906-911.
241. Bastos I, Mallya J, Ingvarsson L, Reimer A, Andreasson L. Middle ear disease and hearing impairment in northern Tanzania. A prevalence study of schoolchildren in the Moshi and Monduli districts. *Int J Pediatr Otorhinolaryngol* 1995; **32**(1): 1-12.
242. Baldwin RL. Effects of otitis media on child development. *Am J Otol* 1993; **14**(6): 601-604.
243. Torre P,3rd, Zeldow B, Hoffman HJ,Buchanan A, Siberry GK, Rice M, et al. Hearing Loss in Perinatally Human Immunodeficiency Virus-Infected and Human Immunodeficiency Virus -Exposed but Uninfected Children and Adolescents. *Pediatr Infect Dis J* 2012; **31**(8): 835-841.
244. Gorga MP, Worthington DW, Reiland JK, Beauchaine KA, Goldgar DE. Some comparisons between auditory brain stem response thresholds, latencies, and the pure-tone audiogram. *Ear Hear* 1985; **6**(2): 105-112.
245. Suchdev P, Ahrens K, Click E, Macklin L, Evangelista D, Graham E. A Model for Sustainable Short-Term International Medical Trips. *Ambul Pediatr* 2007; **7**(4): 317-320.
246. Decamp M. Ethical review of global short-term medical volunteerism. *HEC Forum* 2011; **23**(2): 91-103.
247. Macklin R. Double standards in medical research in developing countries. Cambridge: Cambridge University Press; 2004.
248. Olusanya BO, Luxon LM, Wirz SL. Ethical issues in screening for hearing impairment in newborns in developing countries. *J Med Ethics* 2006; **32**(10): 588-591.
249. McPherson B. Newborn hearing screening in developing countries: needs & new directions. *Indian J Med Res* 2012; **135**: 152-153.
250. Pirozzo S, Papinczak T, Glasziou P. Whispered voice test for screening for hearing impairment in adults and children: systematic review. *BMJ* 2003; **327**(7421) :967.

# APPENDIX

Study forms translated from Portuguese originals.

## HISTORY AND PHYSICAL EXAMINATION

page 1

Name:	Residence:	
Date of birth:	Age:	Telephone:

Study number \_\_\_\_\_

HOSPITAL PEDIÁTRICO DAVID BERNARDINO

Gender Female ☐

Date of admission: \_\_\_\_/\_\_\_\_/\_\_\_\_

Male ☐

Weight : \_\_\_\_ Kg Upper middle arm circumference: \_\_\_\_ cm Height \_\_\_\_ cm

Head circumference: \_\_\_\_ cm Blood pressure \_\_\_\_/\_\_\_\_ mmHg, P \_\_\_\_

### HISTORY

YES NO Unknown

HOUSEHOLD SIZE \_\_\_\_ persons, \_\_\_\_ adults, \_\_\_\_ children

RUNNING WATER AT HOME ☐ ☐ ☐

ELECTRICITY AT HOME ☐ ☐ ☐

BREASTFEEDING Until \_\_\_\_ months ☐ ☐ ☐

SOLID FOOD Introduced at \_\_\_\_ months ☐ ☐ ☐

ATTENDANCE AT VACCINATION PROGRAM ☐ ☐ ☐

ANTES SAUDÁVEL ☐ ☐ ☐

CHRONIC DISEASES ☐ ☐ ☐

PREVIOUS DISEASES \_\_\_\_\_

MALARIA ☐ ☐ ☐

TYPHOID FEVER ☐ ☐ ☐

TUBERCULOSIS ☐ ☐ ☐

HIV / AIDS ☐ ☐ ☐

PREVIOUS ADMISSIONS ☐ ☐ ☐

MEDICATION; WHICH? ☐ ☐ ☐

ANTIBIOTIC; WHICH? ☐ ☐ ☐

QUININE BEFORE ☐ ☐ ☐

SURGERY BEFORE ☐ ☐ ☐

REASON FOR CONSULTATION \_\_\_\_\_

EARACHE; Specify ☐ ☐ ☐

FEELS THAT HEARS WELL; Specify ☐ ☐ ☐

HEARING PROBLEMS IN COMPARISON WITH OTHER CHILDREN ☐ ☐ ☐

HEARING DIFFICULTIES IN THE FAMILY ☐ ☐ ☐

Specify \_\_\_\_\_

### PHYSICAL EXAMINATION

YES NO Unknown

GOOD GENERAL CONDITION: ☐ ☐ ☐

SIGNS OF MALNUTRITION: ☐ ☐ ☐

☐ slight ☐ mod. ☐ grave

FOCUS OF INFECTION Specify ☐ ☐ ☐

HAEMOGLOBIN \_\_\_\_ g/dl

OTHER INFORMATION \_\_\_\_\_

**OTORHINOLARYNGOLOGICAL STATUS**

**YES      NO      Unknown**

FACE NORMAL

☐      ☐      ☐

Specify: \_\_\_\_\_

AURICLES NORMAL

☐      ☐      ☐

Specify: \_\_\_\_\_

NOSE : Direct / Deviated \_\_\_\_\_

NASAL SEPTUM: Direct / Deviated \_\_\_\_\_

RIGHT NOSTRIL: Rhinorrhea / Crust / Polyp / Normal \_\_\_\_\_

LEFT NOSTRIL: Rhinorrhea / Crust / Polyp / Normal \_\_\_\_\_

LIPS-GINGIVAS-PALATE NORMAL \_\_\_\_\_

☐      ☐      ☐

Specify: cleft palate \_\_\_\_\_

DENTITION NORMAL \_\_\_\_\_

☐      ☐      ☐

TEETH: Healthy / Caries / Obturated / Missing \_\_\_\_\_

ORAL MUCOSA NORMAL \_\_\_\_\_

☐      ☐      ☐

Specify: Vesicles / Leukoplakia / Erythroplakia / Candidiasis \_\_\_\_\_

TONSILS NORMAL \_\_\_\_\_

☐      ☐      ☐

Right tonsil: 0, 1, 2, 3, 4+ \_\_\_\_\_ Left tonsil: 0, 1, 2, 3, 4+ \_\_\_\_\_

UVULA NORMAL \_\_\_\_\_

☐      ☐      ☐

TONGUE: Normal / Big \_\_\_\_\_

LINGUAL FRENULUM: Normal / Short \_\_\_\_\_

SPEECH NORMAL \_\_\_\_\_

☐      ☐      ☐

Specify: Not comprehensible / Stuttering / Pronunciation difficulties \_\_\_\_\_

VOICE: Clear / Hoarse \_\_\_\_\_

STRIDOR: Reason? \_\_\_\_\_

☐      ☐      ☐

EYE MOVEMENTS NORMAL \_\_\_\_\_

☐      ☐      ☐

Specify: Right / Left / Strabismus / Nystagmus \_\_\_\_\_

OTHER INFORMATION: \_\_\_\_\_

DOCTOR'S SIGNATURE \_\_\_\_\_



page 3

## OTORHINOLARYNGOLOGICAL STATUS

YES NO UNKNOWN

PALPATION OF NECK NORMAL\_\_\_\_\_

☐
☐
☐

Specify: Tumor / Cyst\_\_\_\_\_

LYMPHADENOPATHY; Lymph nodes, size\_\_\_\_\_cm

☐
☐
☐

FISTULES\_\_\_\_\_

☐
☐
☐

Specify: Right / Left / Preauricular / Type I / Type II / Secretion

☐
☐
☐

SALIVARY GLANDS NORMAL\_\_\_\_\_

PAROTID

Right:

Left:

SUBMANDIBULAR

Right:

Left:

SUBLINGUAL

Right:

Left:

☐
☐
☐

FACIAL NERVE NORMAL\_\_\_\_\_

FROWNING

Dto:

Esq:

CLOSING EYES

Dto:

Esq:

SMILING

Dto:

Esq:

WEBER\_\_\_\_\_

	RIGHT EAR	LEFT EAR
RINNE		
OTORRHEA		
<b>OTOSCOPY:</b> Auditory canal		
<b>OTOSCOPY:</b> _____		
Tympanic membrane		
Mobility of tympanic membrane		
<b>TYMPANOMETRY</b>		
<b>HEARING TESTS</b>		
DATA		
NAME OF EXAMINATOR		
<b>BAEP COMPLETED</b>		
HEARING THRESHOLD, dB		
<b>OAE COMPLETED</b>		

OTHER INFORMATION\_\_\_\_\_